



Journal of Nutrition and Food Security

Shahid Sadoughi University of Medical Sciences
School of Public Health
Department of Nutrition
Nutrition & Food Security Research Center



eISSN: 2476-7425

pISSN: 2476-7417

JNFS 2023; 8(4): 631-638

Website: jnfs.ssu.ac.ir

The Efficacy of Egg White Powder on Anthropometric Indices in Malnourished Children

Maryam Hamidian Shirazi; MSc^{1,2,3}, Afsane Ahmadi; PhD³, Mohammad Reza Dehghani; PhD⁴,
Mohammad Jafar Dehzad; MSc³, Maryam Ekramzadeh; PhD^{*2} & Mehran Nouri; PhD^{1,2}

¹ Student Research Committee, Shiraz University of Medical Sciences, Shiraz, Iran; ² Nutrition Research Center, Shiraz University of Medical Sciences, Shiraz, Iran; ³ Department of Clinical Nutrition, School of Nutrition and Food Sciences, Shiraz University, Shiraz University of Medical Sciences, Shiraz, Iran; ⁴ Clinical Education Research Center, Shiraz University of Medical Sciences, Shiraz, IR Iran.

ARTICLE INFO

ORIGINAL ARTICLE

Article history:

Received: 17 Jan 2022

Revised: 28 Mar 2022

Accepted: 28 Mar 2022

*Corresponding author:

mekramzade@gmail.com

Assistant Professor, Nutrition Research Center, Department of Clinical Nutrition, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Razi Blvd, Shiraz, Iran.

Postal code: 7153675500

Tel: +98 71 37251004

ABSTRACT

Background: White egg powder provides essential amino acids compounds. This study aims to investigate the theory that a regular consumption of white egg powder would improve weight gain and appetite in children with malnutrition. **Methods:** The present experimental study was conducted among children aged 3-6 years with mild to severe malnutrition referred to Ghadir Mother and Child Hospital affiliated with Shiraz University of Medical Sciences from April to October 2017. The malnourished children received egg white protein (1 g/kg of their current body weight) for 3 weeks. Weight loss as a primary outcome and weight-for-height Z-score (WHZ), height-for-age Z-score (HAZ), weight-for-age Z-score (WAZ), and BMI-for-age Z-score (BAZ) were measured as secondary outcomes, assessed at baseline and after six weeks of study. **Results:** Of the 23 children included, 20 finished the study. Weight increased in all children although it was not significant ($P = 0.101$). A lower appetite was seen at the end of the intervention, but it was not significant ($P = 0.575$). Malnutrition based on WHZ and BAZ decreased compared to the baseline measurement, while the changes were not significant. Regarding other markers, no statistically significant changes were found compared to the baseline measurement. **Conclusion:** The addition of white egg powder to complementary nutrition could not improve weight gain and appetite in malnourished children. Further research with a larger study population and longer intervention time is needed to demonstrate the clear effect of egg white on improving malnutrition in children.

Keywords: White egg powder, Malnutrition, Children, Weight gain

Introduction

Malnutrition is a condition in which the body does not receive enough nutrients, leading to acute or chronic deficiencies in protein, other nutrients, and energy in children that can cause

half of the deaths among them (Kemenkes, 2011). Malnutrition can lead to cognitive impairment and growth retardation and is also related to a higher vulnerability to different types of morbidity and

This paper should be cited as: Hamidianshirazi M, Ahmadi A, Dehghani MR, Dehzad MJ, Ekramzadeh M, Nouri M. *The Efficacy of Egg White Powder on Anthropometric Indices in Malnourished Children. Journal of Nutrition and Food Security (JNFS)*, 2023; 8(4): 631-638.

mortality in children (Black *et al.*, 2008). The prevalence of malnutrition in developing countries is due to economic and social problems, which increase the risk factor of physical complications and mortalities in children (Fenton and Silverman, 2004). About 26.9 million children have severe malnutrition in southern Asia (Unicef *et al.*, 2012). According to recent studies, 6 percent of the under-five children in Iran are underweight (Kia *et al.*, 2017). Imbalance between nutrient consumption and metabolic requirements causes symptoms of malnutrition, which, as a consequence, leads to a variety of clinical illnesses (Norman *et al.*, 2008). Adequate and balanced nutrition plays a key role in children's physical and mental growth (World Health Organization, 2006). Research has mainly employed supplements, but inadequate evidence exists for nutrients. Eggs may improve the immune systems of malnourished children (Iannotti *et al.*, 2014). Eggs are rich in choline, betaine, and vitamin B12, and provide all the essential amino acids for a baby's growth (Iannotti *et al.*, 2014). Adequate and high-quality protein in the diet is essential for building and preventing muscle atrophy and also provides adequate energy supply and growth (Leidy *et al.*, 2007). In a malnourished child, branched-chain amino acids affect compensatory growth, protein synthesis, and weight gain (Hsu *et al.*, 2014).

A study showed that the frequency of supplements increased nutrition and weight gain in the intervention group compared to the control group (Roy *et al.*, 2007). In addition, in a clinical trial study, egg consumption in malnourished children resulted in a 9-fold weight gain compared to the control group (Setyaningsih, 2016). The Lulun project aimed to determine the effectiveness of eggs on biomarkers of early childhood nutrition and growth. The results of this study showed that eggs reduced short stature and increased linear growth with a length z score for age (Iannotti *et al.*, 2017). Egg whites improve children's growth, but there are no comprehensive and appropriate studies in this field. Therefore, given the importance of children's development and the importance of time in the treatment of their malnutrition, this clinical

experimental study aimed to assess the efficacy of egg white powder as a supplementary food on the anthropometric status of children with mild to moderate malnutrition.

Materials and Methods

Study design and participants: This 6-week experimental study was conducted on children with mild to severe malnutrition at Ghadir Mother and Child Hospital, affiliated with Shiraz University of Medical Sciences from April to October 2017. The children referred to a pediatrician due to their short stature and low weight and after the doctor's diagnosis, those without underlying disorders were included in the study. Children aged 3-6 years with mild to severe malnutrition and a history of using growth drugs or supplements during the study met eligible criteria. The exclusion criteria included patients who were sick, dying, taking any kind of medicine or supplement, and did not consume egg white protein powder based on the required amount. A sample size of 20 participants was designed based on weight: X1: 6.4, X2: 7.2, SD1: 0.9, and an alpha error probability of 0.05, power = 0.80, according to a similar study (Setyaningsih, 2016). In this experimental study, 23 children were selected from the group as the final sample size. Dietary adherence was measured through subjective methods (self-reported adherence rate, which ranged from 70% to 90%).

Based on the evidence, the Dietary Reference Intakes for protein indicate that children aged 1-3 years and 4-8 years require 1.1 and 0.95 g/kg/day, respectively, based on the classic nitrogen balance technique (Paula *et al.*, 2002). Therefore, the authors believed that the amount of egg white protein required by each child was calculated based on their current weight (1 g/kg of body weight), and parents were given a packet of egg white protein powder (300 g) for three weeks. Parents were asked to add two or three servings of egg white protein powder to yogurt, stews, soups, or other watery foods daily (they were not too hot), and they were asked not to let the children notice.

In addition, the parents were counseled by the dietitian, and they were then trained in

healthy balanced diets and different techniques of decorating baby food to stimulate appetite, as well as a booklet. A daily registration form was given to the parents to record the amount of protein white egg powder consumed by their children, and they were asked to complete it daily for a more accurate assessment of supplement consumption. The package was given to the parents at the beginning of the study and again in the third week. Parents were asked to give back the empty package in the follow-up meetings (at weeks 3 and 6 of the intervention).

Furthermore, researcher reminded the participants to consume egg powder once a week. Their parents were requested not to change the common powder consumption of their children through the experimental study.

Measurements: The weight and height of the participants were evaluated from the beginning to the end of the study based on standard methods (de Onis *et al.*, 2004). Height was measured by a standard tape measure, barefoot, with the child close to the wall and looking straight ahead, with an error of about one-tenth of a centimeter (Seca 220, Hamburg, Germany). Weight was measured using Seca hand scales, with a minimum wear and error of nearly 100 grams (Seca 755, Hamburg, Germany).

The Z-score was used to determine the difference between normal status and malnutrition in children, which was calculated by diagram. Also, demographic questions (name, surname, age, current height, and weight, history of illness or medications taken, and appetite status) were asked from the beginning of the study.

Ethical considerations: The study was conducted in accordance with the Declaration of Helsinki and Good Clinical Practice guidelines and was accepted by the Ethics Committee of Shiraz University of Medical Sciences (IR.SUMS.REC.1396.162). This clinical experimental study was registered in the Iranian Registry of Controlled Experimental Studies (IRCT) (IRCT20100223003408N9). Consent form was obtained from the children's parents.

Data analysis: The data were analyzed using the

SPSS statistical software. The normal distribution of the data was checked by the Kolmogorov-Smirnov test. A paired t-test was used to assess changes in each group and compare them within groups. ANOVA was performed to assess differences in total weight between the growth indicators. The association between weight and growth indicators was analyzed using linear regression model. The significance level was set at $P\text{-value} \leq 0.05$. The categorical variable was described by frequencies and percentages.

Results

Of the 25 children evaluated for eligibility, 2 children were not eligible to enter the study and 3 left the study (**Figure 1**). Except for one case of nausea after taking the powder on the first day, no side effects or problems were reported due to taking the powder. Baseline demographic and clinical parameters are shown in **Table 1**.

Table 1. General children's characteristics at baseline.

Variables	
Gender	
Male	14(61.0)
Female	9(39.0)
Mother's education	
High school	2(8.7)
Associate degree	17(73.9)
Bachelor's degree	3(13.0)
Master's degree	1(4.3)
Place of residence	
Urban	20(87.0)
Rural	3(13.0)
Constipation	5(21.7)
Vomiting	1(4.3)
Anemia	14(60.9)
Low appetite	17(79.3)
Duration of breast feeding (month)	16.26±9.49
Birth weight (kg)	3.03±0.86
Height (cm)	99.40±7.74
Age(month)	48.78±10.56

Values are presented as mean ± SD or n (%).

The prevalence of anemia and anorexia was high in participants at the beginning of the study (**Table 1**). Changes in clinical outcomes during the treatment phase of this study are demonstrated in **Table 2**. The increment in WHZ and BAZ levels in

participants could have some clinical values. It might be indicative of practical effects of white egg powder in patients although these changes were not significant during the study ($P=0.168$ and $P=0.089$, respectively). Appetite levels decreased

during the study, but it was not significant. There were no significant differences in HAZ and other parameters at the beginning and end of the intervention (**Table 2**).




Figure 1. Flow diagram of the experimental study.

Table 2. Changes in growth indicators after 6-week supplementation with white egg powder.

Parameters	Before	After	Changes (CI)	P-value ^a
Weight (kg)	15.18±2.43 ^b	15.27±2.45	0.086±0.24 (-0.18 to 0.19)	0.10
WHZ	-0.32±0.72	-0.26±0.72	0.062±0.19 (-0.28 to 0.15)	0.16
HAZ	-0.62±1.04	-0.62±1.04	0	1
WAZ	-0.54±0.63	-0.53±0.67	0.013±0.13 (-0.40 to 0.70)	0.61
BAZ	-0.27±0.76	-0.19±0.75	0.08±0.2 (-0.01 to 0.17)	0.08
Appetite score	0.74±0.44	0.65±0.48	-0.087±0.73 (-0.40 to 0.23)	0.57

BMI: Body mass index (kg/m²); BAZ: Body mass index z score; HAZ: Height-for-age z score; WAZ: Weight-for-age z score; WHZ: Weight-for-height z score; CI: Confidence interval; ^a: Paired t-test; ^b: Mean± SD

Discussion

The data indicated that supplementation with the weighted egg powder improves growth indicators, including weight, but no statistically significant

difference was found during the study.

Children with malnutrition need to improve their health. They need protein, fat, minerals, and vitamins (Hossain *et al.*, 2020). In a study in this

field, complete milk protein supplementation in school-age children was reported to significantly increase weight and height compared to control group who followed a routine school diet without receiving extra protein from milk (Lampl *et al.*, 1978). A meta-analysis study by Laura Pimpin *et al.* showed that animal protein generally causes weight gain in children (Pimpin *et al.*, 2019). The WHO has considered animal protein as a complementary food in the management of malnutrition in children and states that animal food improves the condition of children by providing amino acids and other nutrient needs (Skau *et al.*, 2015). Consistent with the present study, protein supplements during pregnancy had no benefit in improving malnutrition status or birth outcomes (Tang and Krebs, 2014). Therefore, significant changes may be observed compared to the present results if higher doses of egg white powder were consumed by children with malnutrition for a longer period of time. This increase in weight can have clinical values. It may indicate the effects of egg white consumption on malnourished children. No significant increase in the WHZ of participants was observed after the intervention. Lulun project had a significant effect on increasing weight-for-age z score by 0.61 (95% CI, 0.45–0.77) (Iannotti *et al.*, 2017). However, we did not observe any significant change in the WHZ after the intervention which could be due to the short duration of this study

Also, in children with malnutrition, a slight improvement in the mean z-scores for the two anthropometric indices, including BAZ, was observed after consumption of egg white powder compared to the beginning of the study, but these changes were not significant. In Lulun project, BAZ also increased significantly in the treatment group compared to the control group after consuming eggs in children for 6 months (Iannotti *et al.*, 2017). This difference is probably due to the larger sample size compared to the present study. An observational study in India reported a significant direct association between low egg consumption and stunted growth in children (Aguayo *et al.*, 2016).

It was hypothesized that eggs, as a high-quality food, would be effective in the compensatory development of malnourished children. Eggs contain a lot of choline, a substance that has been shown to increase growth in animal models (Zeisel and Da Costa, 2009).

The effectiveness of animal-source proteins on children's growth has been confirmed (Eaton *et al.*, 2019, Kaimila *et al.*, 2019). Branched-chain amino acids are essential for the growth of children (Nuss and Tanumihardjo, 2011, Pillai and Kurpad, 2012). Observational studies have revealed that consuming fewer eggs resulted in malnutrition in children (Aguayo *et al.*, 2016). Eggs are rich in amino acids and other nutrients that are effective in child growth (Zeisel and Da Costa, 2009). Therefore, high-quality diets like eggs could provide calories and protein and also improve appetite in young children. Consistent with the present study, in a study in rural Malawi, growth was not improved by consuming one egg per day (Iannotti *et al.*, 2017).

The lack of change in anthropometric indicators in the current study could be due to the short duration of intervention and also different health and socioeconomic conditions. Increasing height was found to be related to income status (McGovern *et al.*, 2017). In low-income environments, getting the essential amino acids is not enough (Kampman-van de Hoek *et al.*, 2016). Like many low- and middle-income countries in the world, Iran is one of the countries in short supply (Kia *et al.*, 2017). The critical lack of living facilities, poverty, and unfavorable food for kids might be the reasons why there is malnutrition in children in Iran (Sharifzadeh *et al.*, 2010). Eggs are a complete, accessible and safe food in poor populations compared to fortified complementary food (Iannotti *et al.*, 2014).

The present study closely monitored allergic reactions to eggs during the study and unfortunately one participant was excluded from the study due to allergies. Egg allergy is one of the most common immunoglobulin-mediated food allergies in children although few studies have been conducted in impoverished populations

(Boye, 2012). According to the American Academy of Sciences, supplemental feeding with eggs is considered without the risk of increasing the incidence of allergies (Greer *et al.*, 2008).

In this study, the effect of egg white powder on weight gain and other anthropometric indices in malnourished children was tested for the first time. This experimental study has some limitations. The possibility of no intervention for malnourished children is not scientifically and morally acceptable. This study was designed and conducted without a control group. It would be preferable to look for other inflammatory agents. Feeding was indirectly checked by telephone, so there was uncertainty about the amount of white egg powder consumed by the studied children. With significant protein and micronutrients found in egg whites, it was expected that the white egg powder would improve growth and appetite in children with malnutrition. Studies with longer intervention time appear to be more effective. Another limitation of the study is the low sample size and short duration. Six weeks is too short to observe the changes in primary and secondary outcomes, especially in terms of anthropometric indicators. It is suggested that egg powder with desirable flavors with better reception by children be evaluated in future studies. It was also better to use appetite tests to accurately assess children's appetite.

Conclusions

This experimental study showed that 1 g/kg white egg powder per day in 3-6-year-old children with malnutrition for 6 weeks can affect the WHZ and BAZ parameters to some extent, but these changes were not significant. Egg allergy was also observed in one participant. Further studies with larger sample sizes and lengths of studies are needed to evaluate the effectiveness of egg whites in the development of malnourished children. Since malnutrition is a multifaceted condition, it will definitely necessitate complicated involvement to improve it.

Acknowledgement

Thanks are owed to the Vice Chancellor of

Research Department of Shiraz University of Medical Sciences for their financial support. Also, the authors would like to thank editorial assistance in Research Consulting Center (RCC) of Shiraz University of Medical Sciences.

Conflict of interest

None

Authors' contributions

Research idea and study design: Ekramzadeh M, Ahmadi A, Dehzad MJ, Dehghani MR; data acquisition: Dehzad MJ, Dehghani MR; data analysis/interpretation: Hamidianshirazi M, Ekramzadeh M, Nouri M; statistical analysis: Hamidianshirazi M, Ahmadi A, Nouri M; supervision or mentorship: Ekramzadeh M, AH. Each author contributed to the important intellectual content during manuscript drafting or revision.

Financial support

This study was financially supported by the Vice-chancellor for Research Affairs of Shiraz University of Medical Sciences (grant No. 96-01-84-14813).

Reference

- Aguayo VM, Nair R, Badgaiyan N, Krishna VJM & nutrition c 2016. Determinants of stunting and poor linear growth in children under 2 years of age in India: An in-depth analysis of Maharashtra's comprehensive nutrition survey. *12*: 121-140.
- Black RE, et al. 2008. Maternal and child undernutrition: global and regional exposures and health consequences. *Lanset*. **371 (9608)**: 243-260.
- Boye JI 2012. Food allergies in developing and emerging economies: need for comprehensive data on prevalence rates. *Clinical translational allergy*. **2 (1)**: 1-9.
- de Onis M, Onyango AW, Van den Broeck J, Chumlea WC & Martorell R 2004. Measurement and standardization protocols for anthropometry used in the construction of a new international growth reference. *Food nutrition bulletin*. **25 (1_suppl_1)**: S27-S36.

- Eaton JC, et al.** 2019. Effectiveness of provision of animal-source foods for supporting optimal growth and development in children 6 to 59 months of age. *Cochrane database of systematic reviews.*(2).
- Fenton M & Silverman E** 2004. Medical nutrition therapy for human immunodeficiency virus (HIV) disease. In *krause's food, nutrition & diet therapy* (ed. I. Mahan and S. Escott-stump). USA: WB Saunders Company: Philadelphia.
- Greer F, Sicherer S & Burks A** 2008. American Academy of Pediatrics Committee on Nutrition, American Academy of Pediatrics Section on Allergy and Immunology. Effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, timing of introduction of complementary foods, and hydrolyzed formulas. *Pediatrics.* **121** (1): 183-191.
- Hossain MI, Huq S, Islam MM & Ahmed T** 2020. Acceptability and efficacy of ready-to-use therapeutic food using soy protein isolate in under-5 children suffering from severe acute malnutrition in Bangladesh: a double-blind randomized non-inferiority trial. *European journal of nutrition.* **59** (3): 1149-1161.
- Hsu JW, et al.** 2014. Dietary supplementation with aromatic amino acids increases protein synthesis in children with severe acute malnutrition. *Journal of nutrition.* **144** (5): 660-666.
- Iannotti LL, Lutter CK, Bunn DA & Stewart CP** 2014. Eggs: the uncracked potential for improving maternal and young child nutrition among the world's poor. *Nutrition reviews.* **72** (6): 355-368.
- Iannotti LL, et al.** 2017. Eggs in early complementary feeding and child growth: a randomized controlled trial. *Pediatrics.* **140** (1).
- Kaimila Y, et al.** 2019. Consumption of animal-source protein is associated with improved height-for-age z scores in rural Malawian children aged 12–36 months. *Nutrients.* **11** (2): 480.
- Kampman-van de Hoek E, et al.** 2016. Dietary amino acid deficiency reduces the utilization of amino acids for growth in growing pigs after a period of poor health. *Journal of nutrition.* **146** (1): 51-58.
- Kemenkes R** 2011. Pedoman pelayanan anak gizi buruk, Departemen Kesehatan Direktorat Jenderal Bina Kesehatan Masyarakat, Jakarta.
- Kia AA, Rezapour A, Khosravi A & Abarghouei VA** 2017. Socioeconomic inequality in malnutrition in under-5 children in Iran: evidence from the multiple indicator demographic and health survey, 2010. *Journal of preventive medicine public health.* **50** (3): 201.
- Lampl M, Johnston FE & Malcolm LA** 1978. The effects of protein supplementation on the growth and skeletal maturation of New Guinean school children. *Annals of human biology.* **5** (3): 219-227.
- Leidy HJ, Carnell NS, Mattes RD & Campbell WW** 2007. Higher protein intake preserves lean mass and satiety with weight loss in pre-obese and obese women. *Obesity.* **15** (2): 421-429.
- McGovern ME, Krishna A, Aguayo VM & Subramanian S** 2017. A review of the evidence linking child stunting to economic outcomes. *International journal of epidemiology.* **46** (4): 1171-1191.
- Norman K, Pichard C, Lochs H & Pirlich M** 2008. Prognostic impact of disease-related malnutrition. *Clinical nutrition.* **27** (1): 5-15.
- Nuss ET & Tanumihardjo SA** 2011. Quality protein maize for Africa: closing the protein inadequacy gap in vulnerable populations. *Advances in nutrition.* **2** (3): 217-224.
- Paula T, Sandra S, Allison AY & Mary P** 2002. reference intakes for energy, carbohydrate, fiber, fat, fatty acids Dietary , cholesterol, protein, and amino acids. *Journal of American diet association.* **102** (11): 1621-1630.
- Pillai RR & Kurpad AV** 2012. Amino acid requirements in children and the elderly population. *British journal of nutrition.* **108** (S2): S44-S49.
- Pimpin L, et al.** 2019. Effects of animal protein supplementation of mothers, preterm infants, and term infants on growth outcomes in childhood: a systematic review and meta-analysis of

randomized trials. *American journal of clinical nutrition*. **110** (2): 410-429.

Roy SK, et al. 2007. Prevention of malnutrition among young children in rural Bangladesh by a food-health-care educational intervention: a randomized, controlled trial. *Food nutrition bulletin*. **28** (4): 375-383.

Setyaningsih P 2016. The Effectiveness of Egg Protein to Malnutrition Recovery in Toddler, <https://publikasiilmiah.ums.ac.id/xmlui/handle/11617/6826>.

Sharifzadeh G, Mehrjoofard H & Raghebi S 2010. Prevalence of malnutrition in under 6-year olds in South Khorasan, Iran. *Iranian journal of pediatrics*. **20** (4): 435.

Skau JK, et al. 2015. Effects of animal source food and micronutrient fortification in complementary food products on body composition, iron status, and linear growth: a

randomized trial in Cambodia. *American journal of clinical nutrition*. **101** (4): 742-751.

Tang M & Krebs NF 2014. High protein intake from meat as complementary food increases growth but not adiposity in breastfed infants: a randomized trial. *American journal of clinical nutrition*. **100** (5): 1322-1328.

Unicef, World Health Organization & World Bank Group 2012. Levels and trends in child malnutrition.

World Health Organization 2006. WHO child growth standards: length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: methods and development. World Health Organization.

Zeisel SH & Da Costa K-A 2009. Choline: an essential nutrient for public health. *Nutrition reviews*. **67** (11): 615-623.