



Journal of Nutrition and Food Security

Shahid Sadoughi University of Medical Sciences
School of Public Health
Department of Nutrition
Nutrition & Food Security Research Center



Shahid Sadoughi University of Medical Sciences
School of Public Health

eISSN: 2476-7425

pISSN: 2476-7417

JNFS 2023; 8(3): 493-520

Website: jnfs.ssu.ac.ir

Comparison of the Effect of a Low-Carbohydrate Diet with a Low-Fat Diet on Anthropometric Indices and Body Fat Percentage: A Systematic Review and Meta-Analysis of Randomized Controlled Trials

Mina Darand; PhD^{1,2}, Shirin Hassanizadeh; MSc³, Sepide Talebi; MSc^{4,5}, Zahra Darabi; MSc^{6,7}
Mohammad Bagherniya; PhD⁸, Fatemeh Yaghoobi; PhD⁹, Shahab Alizadeh; PhD¹⁰, Zeinab Darand; MD¹¹,
Yasaman Azamian; MSc¹² & Hadi Abdollahzad; PhD^{*13}

¹ Department of Clinical Nutrition, School of Nutritional Sciences and Dietetics, Isfahan University of Medical Sciences, Isfahan, Iran; ² Nutrition and Food Security Research Center, Isfahan University of Medical Sciences, Isfahan, Iran; ³ Department of Community Nutrition, School of Nutrition and Food Science, Nutrition and Food Security Research Center, Isfahan University of Medical Science, Isfahan, Iran; ⁴ Students' Scientific Research Center (SSRC), Tehran University of Medical Sciences, Tehran, Iran; ⁵ Department of Clinical Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran; ⁶ Department of Nutrition, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran; ⁷ Nutrition and Food Security Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran; ⁸ Nutrition and Food Security Research Center and Department of Community Nutrition, School of Nutrition and Food Science, Isfahan University of Medical Sciences, Isfahan, Iran; ⁹ Esfarayen Faculty of Medical Sciences, Esfarayen, Iran; ¹⁰ Baharloo Hospital, Tehran University of Medical Sciences, Tehran, Iran; ¹¹ School of Veterinary Medicine, University of Kermanshah, Kermanshah, Iran; ¹² Student Research Committee, School of Nutritional Sciences and Food Technology, Kermanshah University of Medical Sciences, Kermanshah, Iran; ¹³ Department of Nutrition, School of Medicine, Urmia University of Medical Sciences, Urmia, Iran.

ARTICLE INFO

SYSTEMATIC REVIEW and META-ANALYSIS

Article history:

Received: 17 Jan 2022

Revised: 26 Feb 2022

Accepted: 19 Mar 2022

*Corresponding author:

hadi_nut@yahoo.com

Department of Nutrition,
School of Medicine, Urmia
University of Medical
Sciences, Serow Highway,
Nazloo, Iran.

Postal code: 5756115111

Tel: +9844 32770698

ABSTRACT

Background: Although many studies have been conducted to compare the effect of adherence to a low-carbohydrate diet (LCD) with a low-fat diet (LFD) on anthropometric indices and body fat percentage, there is still no definite conclusion in this regard. Therefore, the present study aims to summarize results of studies comparing a LFD and a LCD on weight loss. **Methods:** A systematic search of databases including PubMed, Scopus, and Cochran Library was performed up to November 2020. All randomized controlled trials (RCTs) comparing the effect of adherence to a LCD with a LFD on anthropometric indices and body fat percentage were included. Search results were limited to English-language publications. Sixty-three RCTs, including 7660 participants, were selected for the present study. **Results:** Pooled analysis indicated that adherence to LCD was significantly associated with a greater reduction in BMI (SMD = - 0.07, 95% CI: -0.14, -0.001; $P = 0.04$), weight (kg) (SMD = - 0.22, 95% CI: - 0.31, - 0.12; $P \leq 0.001$), and percentage of body fat mass (SMD = - 0.28, 95% CI: -0.48, - 0.08; $P = 0.006$) compared to LFD. However, no significant difference in changes of kilogram of body fat mass and waist circumference was observed between the two diet programs. **Conclusion:** Overall, adhering to LCD was more effective than LFD in losing weight and body fat percentage.

Keywords: Low-fat; Low-carbohydrate; Weight; Obesity

This paper should be cited as: Darand M, Hassanizadeh Sh, Talebi S, Darabi Z, Bagherniya M, Yaghoobi F, et al. Comparison of the Effect of a Low-Carbohydrate Diet with a Low-Fat Diet on Anthropometric Indices and Body Fat Percentage: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Journal of Nutrition and Food Security (JNFS)*, 2023; 8(3): 493-520.

Introduction

Obesity has become one of the most global well-known public health issue, its prevalence has nearly tripled since 1975 (Abarca-Gómez *et al.*, 2017, Zhang *et al.*, 2020). Although several research studies have been conducted to prevent and treat obesity, obesity prevalence is increasing worldwide and it is likely to reach maximum levels between 2026 and 2054 (Janssen *et al.*, 2020).

Lifestyle modification is considered the first line strategy for weight loss and weight maintenance, and diet is the main component of this approach (Wadden *et al.*, 2007, Wadden *et al.*, 2012). Among many diet approaches for weight reduction, diets with altered macronutrient composition such as low-fat diets (LFD) and low-carbohydrate diets (LCD) have been the focus of a large body of evidence. Reduction of dietary fats (low-fat diets were defined to have $\leq 30\%$ total fat content) (Lichtenstein and Van Horn, 1998, Nordmann *et al.*, 2006, van Zuuren *et al.*, 2018) are theoretically reasonable approach for weight loss for several reasons, one them is high calorie density of fats compared with carbohydrates or protein (Buchholz and Schoeller, 2004, Hite *et al.*, 2011, Tobias *et al.*, 2015). Furthermore, studies have examined the association between dietary fat intake and cardiovascular risk and reported that LFD was traditionally applied for weight reduction (Keys *et al.*, 1986, Sackner-Bernstein *et al.*, 2015). However, clinical trials results are inconsistency and evidence does not support LFD superiority to reduce weight and body composition change in comparison to other dietary interventions (Harcombe *et al.*, 2015, Mansoor *et al.*, 2016, Nordmann *et al.*, 2006, Schwingshackl and Hoffmann, 2013, Tobias *et al.*, 2015). On the other hand, LCD characterized by low amounts of carbohydrates and higher quantities of fat and protein is another popular diet for weight loss and control concentrating on macronutrient distribution of the diet (Mansoor *et al.*, 2016). Generally, LCD contains 30-130 g of carbohydrate per day and a very low-carbohydrate ketogenic diet comprises less than 30 g of carbohydrate per day, which usually

permit ketosis to occur (Hite *et al.*, 2011). It is hypothesized that severe carbohydrate restriction results in the depletion of glycogen stores and the excretion of bound water (Astrup *et al.*, 2004). In addition, this diet is an appetite suppressing due to its ketogenic nature, increases satiety, and reduces spontaneous food intake due to its high protein-content. Moreover, the restriction of carbohydrates limits food choices and subsequently reduces energy consumption (Astrup *et al.*, 2004). In a recent meta-analysis, it is shown that adherence to LCD was associated with a decrease in body fat. It is hypothesized that low carbohydrate diets contain high amounts of protein, converting to amino acids in the body. Amino acids increase muscle growth, leading to an increase in muscle mass and subsequently decreasing fat mass (FM) (Hashimoto *et al.*, 2016). On the other hand, in a previous trial, it has been shown that an energy-restricted, high-protein, LFD had no significant impact on body fat in comparison to a high-carbohydrate LFD in obese women (Noakes *et al.*, 2005).

For a long time, debates have existed about the effectiveness of these two kinds of very popular diets in the treatment of obesity and there is still no definite conclusion in this regard particularly in terms of body composition. Therefore, this conclusive systematic review and meta-analysis were undertaken to update previous investigations to assess the effectiveness of LFD interventions in comparison to LCD trials in terms of anthropometry parameters (body mass index (BMI), weight, waist circumference (WC)) as well as body composition (percentage and kilogram of FM).

Materials and Methods

Search strategy: The Preferred Reporting Items for Systematic Reviews and Meta-Analyses Guidelines (PRISMA) were applied for writing the current study (Moher *et al.*, 2015). Scopus, Google Scholar, PubMed, Web of Science and Cochran Library were searched up to November 2020 without any restrictions to identify eligible trials. The effect of a low carbohydrate diet with LFD on anthropometry parameters (BMI, weight, WC) and

body composition (percentage and kilogram of FM) were investigated through Medical Subject Heading (MeSH) terms and non-MeSH terms. Moreover, the references list of the included trials was checked for further possible sources as well as to ensure about the comprehensiveness of searches.

Selection criteria: The selected studies had following criteria: (1) randomized controlled trials (RCTs) design (with any duration and any sample size were included without relevance to physical activity or other factorial interventions), (2) investigating the impact of LCD with LFD on anthropometric indices and body composition (percentage and kilogram of FM) (LCD contains 30-130 g of carbohydrate per day and very low-carbohydrate ketogenic diet comprises less than 30 g of carbohydrate per day, which usually permit ketosis to occur) (Chawla *et al.*, 2020) (these limits were chosen upon consulting literature regarding LCD (Diabetes, 2017), and LFD (Lichtenstein and Van Horn, 1998, Nordmann *et al.*, 2006), and consensus was reached between the authors), (3) reporting macronutrients percentage in the diet, (4) encompassing participants aged ≥ 18 years, and (5) presenting sufficient information linking anthropometric indices and body.

Study selection: Two independent researchers (Darand M and Talebi S) carried out initial screening regarding the articles' titles and abstracts. Then, for selecting RCTs about the effect of a low carbohydrate diet with LFD on anthropometric indices and body fat percentage, full texts of all related articles were assessed by reviewers. Ultimately, any possible disagreement was negotiated and solved via consultation with other researchers (Alizadeh A and Abdollahzad H) (**Figure 1**).

Data extraction: Data extraction was conducted from the selected studies according to criteria including authors' family names, publication year, loss to follow-up, sample size, study duration, diet type, any other intervention, participants' gender, age, and target population, parallel or crossover study design, mean changes and standard deviation (SD) of anthropometric characteristics' levels, as

well as mean and SD of anthropometric indices and body fat percentage at the beginning and the end of the trial (**Table 1**).

Quality assessment: The risk of bias in the included trials was evaluated based on the Cochrane criteria (Higgins *et al.*, 2019). Items for assessing the risk of bias in each research are as follows: (1) random sequence generation; (2) allocation concealment; (3) blinding of participants and personnel; (4) blinding of outcome assessment; (5) incomplete outcome data; (6) selective outcome reporting; and (7) other potential sources of bias. According to the Cochrane Handbook recommendations, three scoring items were applied for studies including "yes", "no" and "unclear" which respectively indicating low risk of bias, high risk of bias, and unclear or unknown risk of bias. Moreover, after determining 'key domains', an overall risk of bias was specified for each trial encompassing good (low risk for all items), fair (low risk for more than three items), and poor (low risk for equivalent and less than three items) (Lorzadeh *et al.*, 2019).

Data synthesis and analysis: The Difference in mean values was defined as effect size. For computing weighted mean differences (WMDs), the mean value was divided by SD of a difference between two random values taken from each group (Higgins, 2011). If the standard error (SE) were reported in studies, SD was obtained by the following equation: $SD = SE \times \sqrt{n}$ (n =number of participants in each group). The WMDs with 95% confidence intervals (CIs) was calculated using a random-effects model for conducting meta-analysis (Borenstein *et al.*, 2011). Cochran's Q test and I-squared (I^2) statistic was applied to determine heterogeneity of trials which was identified by the following criteria: Q statistic P-value of <0.1 ; weak heterogeneity: $I^2 = 25-50$, relatively high heterogeneity: $I^2 = 50-75$, high heterogeneity: $I^2 = 75-100$ (Higgins and Thompson, 2002). Sub-group analysis was performed to identify possible sources of heterogeneity among the selected trials. As the findings of these trials might be affected by intervention duration, sub-group analysis was

accomplished according to these variables. Publication bias was also evaluated through assessing funnel plot and asymmetry tests including Egger's regression test and Begg's rank correlation test at P-value of <0.05 (Duval and Tweedie, 2000). Sensitivity analysis was conducted through individual removal of each study and recalculation of pooled estimates, to determine the impact of a specific trial or a particular group of trials. Meta-regression was also carried out to assess the relationship of estimated effect size with trial duration. Stata software, version 11.2 (Stata Corp.) was used for performing statistical analyses. Statistically significant levels were specified at P-value < 0.05 .

Data analysis: Sensitivity analysis revealed that the overall effects of a LCD on BMI were significantly affected by removing some studies (Brinkworth *et al.*, 2009, de Luis *et al.*, 2007, Frisch *et al.*, 2009, Gardner *et al.*, 2007, Gardner *et al.*, 2018, Haufe *et al.*, 2011, Jang *et al.*, 2017, Lodi *et al.*, 2020, Perna *et al.*, 2019, Phillips *et al.*, 2008, Rajaie *et al.*, 2013, Vander Wal *et al.*, 2007, Wycherley *et al.*, 2010). However, sensitivity analysis did not represent any change in results for weight change, WC, change in FM percentage, and change in FM (kg).

Meta-regression analysis showed that the effect of the LCD versus the LFD on the study outcomes was not modified by the follow-up duration of studies and age of participants (BMI: slope_{Duration}: -0.001; 95 % CI -0.004, 0.001; $P = 0.21$, slope_{Age}: -0.004; 95 % CI -0.01, 0.005; $P = 0.39$), (weight: slope_{Duration}: 0.001; 95 % CI -0.0004, 0.002; $P = 0.14$, slope_{Age}: -0.002; 95 % CI -0.009, 0.003; $P = 0.38$), (Change in percentage of FM: slope_{Duration}: 0.006; 95 % CI -0.0005, 0.01; $P = 0.06$, slope_{Age}: -0.01; 95 % CI -0.03, 0.009; $P = 0.28$).

Publication bias: Publication bias was not observed based on funnel plots and asymmetry tests (**Table 2, Figure 7**)

Ethical consideration: The present study was approved by the Ethical Committee of Kermanshah University of Medical Sciences

Results

Study characteristics: The initial literature search identified a total of 2514 publications. The flow chart reporting the study selection process is presented in **Figure 1**. A total of 327 duplicate studies were first removed and 2077 unrelated records were excluded based on titles and abstracts. Then, 110 articles underwent full-text screening. For studies that reported results for different groups in stratified analysis, all appropriate data were extracted and compared with the control group. Accordingly, two data sets were extracted for studies conducted by Gardner *et al.* (Gardner *et al.*, 2007), Rock *et al.* (Rock *et al.*, 2016), Burgess *et al.* (Burgess *et al.*, 2017), and Petrisko *et al.* (Petrisko *et al.*, 2020). These studies were conducted on two groups of participants with different health status. Finally, 63 relevant RCT studies (67 data sets), with a total sample size of 7660 participants (4434 for the low-carbohydrate arm and 3266 for the low-fat arm), published between 1995 and 2020, were eligible for this meta-analysis according to the inclusion criteria. The obtained data included 30 studies on BMI, 63 studies with 67 data sets on weight, 21 studies with 22 data sets on WC, 8 studies on the percentage of body FM, and 13 studies with 14 data sets on kilogram of body FM. The sample size of the analyzed studies was between 4 to 648 individuals and the duration of follow-up ranged from 2 to 144 weeks. Of the included studies, 24 data sets were on overweight/obese patients with obesity-related comorbidities and the remaining studies were on apparently healthy overweight/obese individuals. Moreover, 14 studies included only women, 1 study included only men (Sharman and Volek, 2004), and other studies included both genders. The studies quality ranged from poor to high quality. The characteristics of the analyzed studies are reported in **Table 1**.

Overall analysis of pooled data: The overall analysis of the included studies revealed that adherence to LCD was significantly associated with a greater reduction in BMI compared to LFD (SMD = -0.07, 95% CI: -0.14, -0.001; $P = 0.04$, **Figure 2**),

weight (SMD = - 0.22, 95% CI: - 0.31, - 0.12; $P \leq 0.001$, **Figure 3**), and percentage of body FM (SMD = - 0.28, 95% CI: -0.48, - 0.08; $P = 0.006$, **Figure 4**). However, no significant difference in changes in kilogram of body FM (**Figure 5**) and WC (**Figure 6**) was observed between the two diet programs.

An additional analysis was performed only on parallel studies which showed similar results. (BMI: SMD = - 0.13, 95% CI: -0.21, -0.05; $P = 0.002$), (weight: SMD = - 0.21, 95% CI: -0.31,-0.11; $P = \leq 0.001$), (WC: SMD = - 0.05, 95% CI: -0.13, 0.02; $P = 0.18$), and (changes of kilogram of body FM: SMD = - 0.04, 95% CI: -0.22, 0.13; $P = 0.61$).

Evidence of heterogeneity was found among the studies comparing the effect of a LCD with a LFD on weight and percentage of body FM. However, no significant heterogeneity was observed among the studies which evaluated BMI, WC, and kilogram of body FM (**Table 2**).

The results of the subgroup analysis are presented in **Table 2**. In subgroup analysis by the duration of follow-up, gender, age, carbohydrate percentage, and health status of participants, a significant BMI lowering LCD effect was supported by studies with ≥ 6 months follow-up (SMD = - 0.14, 95% CI: -

0.23 to - 0.04; $P = 0.004$), age less than 60 years (SMD = - 0.08, 95% CI: - 0.16 to - 0.002; $P = 0.04$), and very low carbohydrate diet (SMD = - 0.19, 95% CI: - 0.31 to - 0.07; $P = 0.001$). However, for weight, this effect was supported by all subgroups, except for overweight or obese individuals with comorbidities, individuals over the age of 60, and diets with more than 10% carbohydrates. In all subgroups, there was no significant difference in WC, except for overweight or obese individuals without comorbidities (SMD=-0.08, 95% CI: - 0.16 to - 0.03; $P = 0.04$). Moreover, for body FM percentage, the mentioned decrease was observed in subgroups of overweight/obese subjects without obesity-related comorbidities (SMD = - 0.21, 95% CI: - 0.37 to - 0.05; $P = 0.009$), after ≥ 6 months follow-up (SMD = - 0.29, 95% CI: - 0.52 to - 0.05; $P = 0.01$), in studies with less than 10% carbohydrates (SMD = - 0.36, 95% CI: - 0.62 to - 0.10; $P = 0.007$), and studies on both genders (SMD = - 0.31, 95% CI: - 0.57 to - 0.05; $P=0.01$). In all subgroup analyses, FM changes (kg) were not significant between the two diet programs (**Table 2**).




Figure 1. Flow chart of studies selection process.




Figure 2. Forest plot illustrating standardized mean difference and 95% confidence intervals for the low-carbohydrate diet versus the low-fat diet on body mass index.




Figure 3. Forest plot illustrating standardized mean difference and 95% confidence intervals for the low-carbohydrate diet versus the low-fat diet on weight.




Figure 4. Forest plot illustrating standardized mean difference and 95% confidence intervals for the low-carbohydrate diet versus the low-fat diet on percentage of fat mass.




Figure 5. Forest plot illustrating standardized mean difference and 95% confidence intervals for the low-carbohydrate diet versus the low-fat diet on kilogram fat mass.




Figure 6. Forest plot illustrating standardized mean difference and 95% confidence intervals for the low-carbohydrate diet versus the low-fat diet on waist circumferences.




Figure 7. Publication bias for outcomes.

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
Struik (Australia; 2020)	Parallel	Obese	Both	59	59	41/43	16	Cho: 14% Pro: 28% Fat: 58% Energy-restricted	Cho: 53% Pro: 17% Fat: 30% Energy-matched energy-restricted	ND	Weight loss
Lodi (Italy; 2020)	Parallel	Overweight	Women	20-35	20- 35	17/17	1.5	Ketogenic diets Cho: 7% Pro: 29% Fat: 64%	Mediterranean diet Cho: 60% Pro: 20% Fat: 20%	ND	Weight loss, BMI- change, body fat
Maekawa (Japan; 2020)	Parallel	Obese	Both	45.6	46.5	18/13	48	Cho: <120 g/d low Cho in combination with IGB therapy	Cho: 50-60% Pro: 15-20% Fat: 20-25% calorie-restricted diet in combination with IGB therapy	ND	Weight loss, BMI- change
Petrisko (American; 2020)	Cross- over	Obese	Both	43.2	43.2	17/17	4	Cho: 10% Pro: 30% Fat: 60% Restricted diet included more animal foods	Cho: 61% Pro: 18% Fat: 21%	1600 and 2200 kcal/d	Weight loss, body fat
Petrisko (American; 2020)	Cross- over	Obese	Both	43.2	43.2	17/17	4	Cho: 10% Pro: 40% Fat: 60%	Cho: 61% Pro: 18% Fat: 21%	1600 and 2200 kcal/d	Weight loss, body fat

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
								Restricted diet included more plant foods and mushrooms			
Perna (Bahrain; 2019)	Parallel	T2DM	Both	59.50	67.7 8	8/9	12	Cho: 27-31% Pro: 22% Fat: 46-50%	Cho: 55-60% Pro: 15-20% Fat: 25-30%	1600-1800 kcal/d	Weight loss, BMI- change, WC change
Guo (American; 2019)	Parallel	Overweight	Both	40.2	39.3	209/205	12/24/48	Cho: 20 g/d	Fat: 20 g/d	~500-600 kcal/d	Weight loss
Gardner (American; 2018)	Parallel	Overweight	Both	40.2	39.3	305/304	48	Cho: 20 g/d	Fat: 20 g/d	ND	Weight loss, BMI- change, WC change, body fat
Burgess (American; 2017)	Parallel	Non-taster	Women	45.3	48.4	26/31	24	Cho: 50 g/d	Fat: 40 to 50 g/d (1,200 to 1,500 kcal/d)	ND	Weight loss
Burgess (American; 2017)	Parallel	Supertaster	Women	47.8	42.1	31/29	24	Cho: 50 g/d	Fat: 40 to 50 g/d (1,200 to 1,500 kcal/d)	ND	Weight loss
Gardner (American; 2016)	Parallel	Pre-menopausal with Insulin resistant	Both	42	44	16/15	24	Cho: 21% Pro: 26% Fat: 53%	Cho: 58% Pro: 20% Fat: 22%	ND	Weight loss

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
Gardner (American; 2016)	Parallel	Pre-menopausal with Insulin sensitive	Both	43	41	14/16	24	Cho: 21% Pro: 20-25% Fat: 20-25%	Cho: 58% Pro: 20% Fat: 22%	ND	Weight loss
Jang (Korea; 2018)	Parallel	NAFLD	Both	ND	ND	52/54	8	Cho: 50-60% Pro: 26% Fat: 53%	Cho: 60-70% Pro: 15-20% Fat: 15-20%	25 kcal/kg IBW	Weight loss, BMI- change, WC change
Tay (Australia; 2018)	Parallel	T2DM	Both	58	58	58/57	96	Cho: 14% Pro: 28% Fat: 58% 35% monounsaturated 13% polyunsaturated fat	Cho: 53% Pro: 17% Fat: 30% 15% monounsaturated 9% polyunsaturated fat	Energy-matched, hypocaloric diets and aerobic/resistance exercise	Weight loss, BMI- change, WC change, body fat
Wycherley (Australia; 2016)	Parallel	Obese and T2DM	Both	58.5	58.4	58/57	52	Cho: 14% Pro: 28% Fat: 58% <10% saturated fat	Cho: 53% Pro: 17% Fat: 30% <10% saturated fat	isocaloric, energy reduced diets (~6 - ~7 MJ/ d) and exercise program	Weight loss
Raygan (Iran; 2016)	Parallel	T2DM	Both	61.1	65.2	28/28	8	Cho: 43-49% Pro: 14-18% Fat: 36-40%	Cho: 60-65% Pro: 14-18% Fat: 20-25%	ND	Weight loss, BMI- change

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
Rock (American; 2016)	Parallel	Obese with Insulin Sensitive	Women	50	50	39/39	48	Cho: 45%	Fat: 20%	ND	Weight loss, WC change
Rock (American; 2016)	Parallel	Obese with Insulin Resistant	Women	50	50	42/43	48	Cho: 45%	Fat: 20%	ND	Weight loss, WC change
Heggen (Norway; 2016)	Parallel	Overweight or obese smokers	Both	49.5	50.8	64/57	4/12	Cho: ≤20% Pro: 25% Fat: 55%	Cho: 50% Pro: 20% Fat: ≤30%	- 500 kcal/d	Weight loss, WC change, body fat
Hu (American; 2015)	Parallel	Obese	Both	45.8	47.8	75/73	12/24/48	Cho: <40 g/d	Fat: <30% < 7 saturated fat	ND	Weight loss
Youssef (Qatar; 2015)	Parallel	Overweight	Women	20-22	20- 22	6/6	6	Cho: 30% Pro: 30% Fat: 40%	Cho: 55% Pro: 30% Fat: 15%	Energy-restricted	Weight loss, BMI- change, WC change, body fat
Bazzano (American; 2014)	Parallel	Obese	Both	45.8	47.8	75/73	12/24/36	Cho: <40 g/d	Cho: 55% Pro: ~15 Fat: <30%	ND	Weight loss, WC change, body fat
Rajaie (Iran; 2013)	Cross- over	Metabolic Syndrome	Both	42.4	42.4	30/30	6	Cho: 43-47% Pro: 15-17% Fat: 36-40%	Cho: 60-65% Pro: 15-17% Fat: 20-25%	ND	Weight loss, BMI- change
Guldbrand (Sweden; 2012)	Parallel	T2DM	Both	61.2	62.7	30/31	24/48/96	Cho: 20% Pro: 30%	Cho: 55-60% Pro: 10-15%	ND	Weight loss, BMI-

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
								Fat: 50%	Fat: 30% < 10% saturated fat		change, WC change
Heggen (Norway; 2012)	Parallel	Obesity with at least one additional metabolic syndrome risk factor	Both	50.3	49.8	78/80	12	Low-Glycemic Load Diet Cho: 30-35% Pro: 25-30% Fat: 35-40%	Cho: 55-60% Pro: 10-15% Fat: <30%	Hypocaloric diets (- 500 kcal/d)	Weight loss
Haufe (Germany; 2011)	Parallel	Obese	Both	43.5	45	84/86	24	Cho: <90 g/d	Cho: <30%	-30% of energy intake before diet	Weight loss, BMI- change, WC change
Foster (American; 2010)	Parallel	Obese	Both	46.2	44.9	153/154	12/24/48/96	Cho: 20 g/d	Cho: 55% Pro: 15% Fat: 30% 1200 to 1800 kcal/d	ND	Weight loss
Klemsdal (Norway; 2010)	Parallel	least one criterion of metabolic syndrome	Both	50.1	49.9	100/102	12/24/48	Cho: 30-35% Pro: 25-30% Fat: 35-40%	Cho: 55-60% Pro: 15% Fat: <30%	ND	Weight loss, WC change
Lim (Australia; 2010)	Parallel	one additional cardiovascula r risk factor	Both	48.3	48.6	27/28	12/60	Cho: 4% Pro: 35% Fat: 60% 20% saturated fat	Cho: 70% Pro: 20% Fat: 10% 3% saturated fat	6500 KJ	Weight loss

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
Mueller (American; 2010)	Parallel	Obese	Both	49	46	9/7	10/20	Cho: 30% Pro: 20% Fat: 50%	Cho: 50% Pro: 20% Fat: 30%	-500 to -750 calorie/d	Weight loss, BMI- change
Thomson (American; 2010)	Parallel	Overweight breast cancer survivors	Women	57.8	57.8	21/19	6/12/18/24	Cho: 35% Pro: 25-30% Fat: 35-40%	Cho: 55-60% Pro: 15-20% Fat: 25%	ND	Weight loss, BMI- change, WC change, body fat
Wycherley (Australia; 2010)	Parallel	Overweight/ Obese	Both	49.9	50.2	26/23	52	Cho: 4% Pro: 35% Fat: 61% 20% saturated fat	Cho: 46% Pro: 24% Fat: 30% <8% saturated fat	isocaloric with moderate energy restriction (~6000 to ~7000 Kj/d)	Weight loss, BMI- change
Yancy (American; 2010)	Parallel	Overweight/ Obese	Both	52.9	52	72/74	48	ketogenic diet Cho: <20 g/d	Fat: <30% <10% saturated fat and <300 mg cholesterol + Orlistat therapy	-500 to 1000	Weight loss, WC change
Bradley (American; 2009)	Parallel	Obese	Both	37.1	40.5	12/12	8	Cho: 20% Pro: 20% Fat: 60%	Cho: 60% Pro: 20% Fat: 20%	ND	Weight loss, BMI- change, WC change, body fat
Frisch (American; 2009)	Parallel	Obese	Both	47	47	100/100	24/48	Cho: <40% Pro: 25% Fat: 35%	Cho: >55% Pro: 15% Fat: 30%	energy-restricted diets	Weight loss, BMI- change,

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
											WC change, body fat
Brinkworth (Australian; 2009)	Parallel	Obesity with at least one additional metabolic syndrome risk factor	Both	51.5	51.4	33/36	8/52	Cho: 4% Pro: 35% Fat: 61%	Cho: 46% Pro: 24% Fat: 30%	isocaloric with moderate energy restriction	Weight loss, BMI- change, body fat
Davis (American; 2009)	Parallel	T2DM	Both	54	53	55/50	12/24/48	Cho: 20-25 g/d	Fat: 25%	ND	Weight loss
Rodriguez- Hernandez (Germany; 2009)	Parallel	Obese	Women	45.3	45.4	21/29	24	Cho: 45% Pro: 27% Fat: 28%	Cho: 54% Pro: 25% Fat: 21% 10% saturated fat	ND	Weight loss, BMI- change, WC change
Iqbal (American; 2010)	Parallel	Obese, Diabetic	Both	60	60	70/74	24/48/96	Cho: 30%	Fat: ≤30% -500 kcal/d	ND	Weight loss
Phillips (American; 2008)	Parallel	Obese	Both	33	38	10/10	4/6	Atkins diet Cho: 20 g/d	Fat: 30%	ND	Weight loss, BMI- change
Stoernell (American; 2008)	Parallel	Hypertriglyce ridemic/ Obese	Both	57	48.4	10/13	8	Atkins diet Cho: 15% Pro: 20-30% Fat: 55-65% <10% saturated fat	Cho: 50-60% Pro: 15% Fat: 30% <10% saturated fat	ND	Weight loss

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
Sloth (Denmark; 2009)	Parallel	Overweight	Both	30	28	15/18	24	Fat: 35-45% high in MUFA, 20%	Fat: 20-30%	ND	Weight loss, BMI- change, body fat
Tay (Australia; 2008)	Parallel	Obese	Both	50.3	51	45/43	24	Cho: 4% Pro: 35% Fat: 61% 20% saturated fat	Cho: 46% Pro: 24% Fat: 30% <8% saturated fat	Isocaloric, (restriction of ~6,000 ~7,000 kJ)	Weight loss
Gardner (American; 2007)	Parallel	Overweight Premenopausal	Women	42	42	77/76	8/24/48	Atkins diet Cho: 20 g/d	Ornish diet Fat: 10%	ND	Weight loss, BMI- change, body fat
DeLuis (Spanish; 2007)	Parallel	Obese	Both	43	42.1	43/47	12	Cho: 38% Pro: 26% Fat: 36%	Cho: 52% Pro: 20% Fat: 27%	1500 kcal/day	Weight loss, BMI- change, WC change, body fat
Ebbeling (American; 2007)	Parallel	Obese	Both	28.2	26.9	36/37	24/72	Low-Glycemic Load Diet Cho: 40% Pro: 25% Fat: 35%	Cho: 55% Pro: 25% Fat: 20%	ND	Weight loss, body fat
LeCheminan (American; 2007)	Parallel	Overweight/ Obese	Both	47.9	45.7	29/26	12/24	Cho: 20%	Fat: 30%	ND	WC change

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
Vanderwal (American; 2007)	Parallel	Overweight	Both	50.46	49.5 8	41/40	4	Cho: 17 g/d Fat: 18 g/d	Cho: 22 g/d Fat: 3 g/d	ND	Weight loss, BMI- change, WC change, body fat
Cardillo (American; 2006)	Parallel	Obese	Both	54	55	27/26	144	Cho: <30 g/d	Fat: <30% - 500 calories	ND	Weight loss
Westman (American; 2006)	Parallel	Obese with LDL-C >130 mg/dl or triglyceride >200 mg/dl	Both	44.2	45.6	59/60	24	ketogenic diets with fish, borage and flaxseed oil supplementation Cho: <20 g/d	Fat: ? reduced-calorie diet	ND	Weight loss
Nickols- richardson (American; 2005)	Parallel	Overweight Premenopaus al	Women	38.8	40.1	13/15	6	Atkins Cho: ≤20 to 40 1500-1700 kcal/d	Cho: 60% Pro: 15% Fat: 25%	ND	Weight loss, BMI- change
Löfgren (Sweden; 2005)	Parallel	Obese	Women	35.7	36.1	20/20	10	Cho: 40-45% Pro: 15-20% Fat: 40-45%	Cho: 60-65% Pro: 15-20% Fat: 20-25%	- 600 kcal/day	Weight loss, BMI- change
Petersen (Germany; 2006)	Parallel	Obese	Both	37.5	37.5	312/336	10	Cho: 40-45% Pro: 15% Fat: 40-45%	Cho: 60-65% Pro: 15% Fat: 20-25%	- 600 kcal/day	Weight loss, BMI- change, WC change, body fat

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
Meckling (Canada; 2004)	Parallel	Obese	Both	41.2	43.2	20/20	10	Cho: 15.4%	Fat: 17.8%	Females: 5020– 6690 and males 5860–9200 kJ/d	Weight loss, BMI- change, body fat
Segal-Isaacson (American; 2004)	Cross- over	Overweight /Obese Premenopausal	Women	52.3	52.3	4/4	12	Cho: 5% Pro: 30% Fat: 65%	Cho: 50% Pro: 30% Fat: 20%	- 200 kcal/d	Weight loss
Sharman (American; 2004)	Cross- over	Overweight	Male	33.2	33.2	15/15	6	Atkins Cho: 10% Pro: 30% Fat: 60%	Cho: 55% Pro: 20% Fat: 25% <10% saturated fat and 300 mg cholesterol	ND	Weight loss
Viguerie (American; 2005)	Parallel	Obese	Both	21-49	21- 49	25/25	10	Cho: 45-50% Pro: 15% Fat: 40-45%	Cho: 60-65% Pro: 15% Fat: 20-25%	Energy-restricted diets	Weight loss, BMI- change, body fat
Yancy (American; 2004)	Parallel	Overweight, Hyperlipidemic	Both	44.2	45.6	59/60	24	Cho: <20 g/d	Fat: <30% <10% saturated fat and <300 mg cholesterol -500 to 1000 kcal	ND	Weight loss, body fat
Volek (American; 2004)	Parallel	Overweight	Women	34	34	13/13	4	Cho: 10% Pro: 30% Fat: 60%	Cho: 55% Pro: 20% Fat: 25%	-500 kcal/d	Weight loss, body fat mass

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
									<10% saturated fat and <300 mg cholesterol		
Brehm (American; 2003)	Parallel	Obese	Women	44.22	43.1 0	22/20	12/24	Cho: 20 g/d (After 2wk of dieting, increase intake of Cho to 40–60 g)	Cho: 55% Pro: 15% Fat: 20%	ND	Weight loss, body fat
Samaha (American; 2003)	Parallel	Obese	Both	53	54	64/68	24	Cho: <30 g/d	Fat: <30% - 500 calories	ND	Weight loss
Petersen (American; 1995)	Crossover	Obese	Women	38.3	38.3	12/12	6	Cho:40%	Cho:55%	ND	Weight loss
Raccette (American; 1995)	Parallel	Obese	Women	41	37	6/7	12	Cho:25% Pro: 25% Fat: 50%	Cho: 60% Pro: 25% Fat: 15%	-5.00 ± 0.56 MJ/d	Weight loss
Foster GD (American; 2003)	Parallel	Obese	Both	44	44.2	20/17	48	Cho: 20 g/d (After 2wk of dieting, increase intake of Cho)	Cho: 60% Pro: 25% Fat: 15%	ND	Weight loss
Shai I (Israel; 2008)	Parallel	Obese	Both	52	51	109/104	96	Cho: 20 g/d (After 2-month of dieting, increase intake of Cho to 120g/d)	Fat: 30%	Mediterranean diet	Weight loss
Dansinger ML	Parallel	Obese or	Both	47	49	21/20	48	Atkins diet	Ornish diet	Weight	Weight

Table 1. Characteristics of eligible studies.

First author (location; year)	RCT design	Target population	Sex	Mean age (years)		Sample size (low CHO diet / low FAT diet)	Duration (Weeks)	Diet Type		Any other intervention	Outcomes
				low CHO diet	low FAT diet			Percentage of macronutrients in low CHO diet	Percentage of macronutrients in low FAT diet		
(American;2007)		overweight						Cho: 20 g/d	Fat: 10%	Watchers, and Zone Diets	loss, BMI- change, WC change

CHO, carbohydrate; PRO, protein; FAT, lipid; T2DM, type 2 diabetes mellitus; NAFLD, nonalcoholic fatty liver disease; IGB, Intra-gastric balloon; IBW, ideal body weight; BMI, body mass index; WC, waist circumference; ND, non-defined; gr; gram, mg; milligram, d; day.

Table 2. Main analyses and prespecified subgroup analyses for effect of low-carbohydrate diet compared with low-fat diet on changes in obesity measures.

Parameters of obesity	Subgroup	Number of datasets	Test of association			Test of heterogeneity		Publication bias		
			SMD	95%CI	P	I ² (%)	P	Begg test	Egger test	
Change in BMI (kg/m ²)	Overall	30	-0.07	-0.14,-0.001	0.04	1.53	0.44	0.17	0.65	
	Follow-up Duration	< 6 months	16	0.01	-0.09, 0.12	0.78	0.0	0.5		
		≥ 6 months	14	-0.14	-0.23,- 0.04	0.004	0.0	0.63		
	Health status	Overweight/obesity without comorbidities	22	-0.06	-0.14, 0.02	0.17	5.63	0.38		
		Overweight/obesity with comorbidities	8	-0.15	-0.32, 0.01	0.07	0.0	0.53		
	Sex	Female	7	-0.16	-0.37, 0.03	0.11	3.86	0.39		
		Both	23	-0.05	-0.13, 0.02	0.15	0.19	0.45		
	% Carbohydrates	LCD	21	-0.01	-0.09,0.08	0.98	0.00	0.65		
		> 50gr/d or 10% VLCD	9	-0.19	-0.31, -0.07	0.001	0.00	0.66		
	Age	≤ 50gr/d or 10%	27	-0.08	-0.16, -0.002	0.04	9.02	0.33		
<60 years ≥60years		3	-0.03	-0.37-0.30	0.83	0.00	0.66			
Change in weight (kg)	Overall	67	-0.22	-0.31, - 0.12	<0.001	67.85	<0.001	0.11	0.05	
	Follow-up Duration	< 6 months	27	-0.28	-0.46, - 0.10	0.002	64.69	<0.001		
		≥ 6 months	40	-0.19	-0.30, - 0.07	0.001	70.43	<0.001		
	Health status	Overweight/obesity without comorbidities	43	-0.30	-0.44, - 0.17	<0.001	74.36	<0.001		
		Overweight/obesity with comorbidities	24	-0.08	-0.20, 0.02	0.13	34.53	0.05		
	Sex	Male	1	-0.69	-1.43, 0.04	0.06	0.00	0.1		
		Female	16	-0.36	-0.73, - 0.002	0.04	82.28	<0.001		
	% Carbohydrates	Both	50	-0.20	-0.29, - 0.11	<0.001	58.46	<0.001		
		LCD > 50gr/d or 10%	39	-0.12	-0.24,0.001	0.05	55.49	<0.001		

Table 2. Main analyses and prespecified subgroup analyses for effect of low-carbohydrate diet compared with low-fat diet on changes in obesity measures.

Parameters of obesity	Subgroup	Number of datasets	Test of association			Test of heterogeneity		Publication bias	
			SMD	95%CI	P	I ² (%)	P	Begg test	Egger test
Change in WC (cm)	VLCD	28	-	-0.5, -0.17	0.00	76.81	<0.001	0.18	0.74
	≤ 50gr/d or 10%	63	0.034	-0.33,-0.12	<0.001	69.87	<0.001		
	Age <60 years	4	-0.07	-0.31,0.15	0.51	0.00	0.76		
	≥60years	22	-0.04	-0.11, 0.03	0.26	0.00	0.46		
	Overall	8	-	-0.12, 0.11	0.95	0.0	0.99		
	Follow-up Duration < 6 months	14	0.003	-0.16, 0.06	0.38	32.84	0.11		
	≥ 6 months	14	-0.08	-0.16, -.003	0.04	0.00	0.94		
	Health status Overweight/obesity without comorbidities	8	0.10	-0.08, 0.28	0.27	30.25	0.18		
	Overweight/obesity with comorbidities	5	0.11	-0.23, 0.46	0.53	48.20	0.1		
	Sex Female	17	-0.05	-0.12, 0.01	0.13	0.0	0.79		
Both	17	-0.01	-0.13,0.10	0.77	17.54	0.25			
% Carbohydrates LCD	5	-0.10	-0.22,0,01	0.09	0.00	0.9			
> 50gr/d or 10%	20	-0.05	-0.14,0.03	0.21	9.75	0.33			
VLCD	2	0.04	-0.4,0.48	0.85	0.00	0.72			
≤ 50gr/d or 10%	8	-0.28	-0.48, - 0.08	0.006	57.38	0.02	0.62	0.36	
Age <60 years	2	-0.29	-0.67, 0.09	0.13	0.0	0.34			
≥60years	6	-0.29	-0.52, - 0.05	0.01	67.49	0.009			
Overall	6	-0.21	-0.37, - 0.05	0.009	25.03	0.24			
Follow-up Duration < 6 months	2	-0.39	-1.18, 0.39	0.32	81.75	0.01			
≥ 6 months	6	-0.21	-0.37, - 0.05	0.009	25.03	0.24			
Health status Overweight/obesity without comorbidities	2	-0.39	-1.18, 0.39	0.32	81.75	0.01			
Overweight/obesity with comorbidities	6	-0.21	-0.37, - 0.05	0.009	25.03	0.24			

Table 2. Main analyses and prespecified subgroup analyses for effect of low-carbohydrate diet compared with low-fat diet on changes in obesity measures.

Parameters of obesity	Subgroup	Number of datasets	Test of association			Test of heterogeneity		Publication bias		
			SMD	95%CI	P	I ² (%)	P	Begg test	Egger test	
Sex	Female	2	-0.21	-0.52, 0.09	0.17	13.53	0.28			
	Both	6	-0.31	-0.57, -0.05	0.01	67.22	0.009			
	LCD	3	-0.07	-0.39, 0.25	0.65	0.00	0.79			
	> 50gr/d or 10%									
% Carbohydrates	VLCD	8	-0.36	-0.62, -0.1	0.007	73.56	0.004			
	≤ 50gr/d or 10%									
Change in fat mass (kg)	Overall	14	0.006	-0.13, 0.14	0.93	24.66	0.18	0.78	0.44	
	Follow-up Duration	< 6 months	9	0.11	-0.004, 0.23	0.05	0.00	0.96		
		≥ 6 months	5	-0.19	-0.48, 0.09	0.19	51.14	0.08		
	Health status	Overweight/obesity without comorbidities	12	0.01	-0.14, 0.17	0.82	31.01	0.14		
		Overweight/obesity with comorbidities	2	-0.06	-0.35, 0.22	0.66	0.0	0.35		
	Sex	Female	3	-0.38	-1.04, 0.27	0.25	53.59	0.11		
		Both	11	0.05	-0.04, 0.16	0.28	0.00	0.60		
	% Carbohydrates	LCD	9	0.02	-0.13, 0.17	0.76	0.00	0.54		
		> 50gr/d or 10%								
		VLCD	5	-0.38	-0.88, 0.12	0.13	54.54	0.11		
≤ 50gr/d or 10%										

Discussion

This is a meta-analysis of RCTs investigating the effect of LCD compared to LFD on anthropometric characteristics. Treatment strategies are important in improving anthropometric indices. Abnormalities of anthropometric characteristics were strongly associated with diseases such as diabetes, hypertension, and cardiovascular disease (Furtado *et al.*, 2018, Hadaegh *et al.*, 2009, Khader *et al.*, 2019). Studies have reported that different diets could affect anthropometric indices. The present meta-analysis study showed that adherence to LCD compared to LFD significantly reduced BMI in studies with ≥ 6 months follow-up, very low carbohydrate diet, and age less than 60 years. The protective effect of LCD on weight was observed except for overweight or obese individuals with comorbidities, individuals over the age of 60 and with diets more than 10% carbohydrates.

A possible reason was that older people generally had less physical activity compared to young people, so that activity-related energy expenditure was also lower. This can lead to increased BMI and weight (Elmadfa and Meyer, 2008). Also, adherence to LCD can reduce FM percentage compared to LFD, but body FM (kg) and WC did not change.

A meta-analysis was similar to the present study published in 2020, investigating the effect of LFD and LCD on weight and lipid profile. In this study, only 38 trials were included and other anthropometric indices were not investigated (Chawla *et al.*, 2020). Previous meta-analysis studies have reported the protective effect of LCD. Another meta-analysis study was conducted on RCTs. It was reported that LCD (50 g carbohydrate or 10% calorie from carbohydrates) can reduce body weight (BW) and FM compared to LCD (about 40% calorie from carbohydrate) (Hashimoto *et al.*, 2016). Another meta-analysis investigated the effect of very-low-carbohydrate ketogenic diets (VLCKD) compared to LFD. Results of this study showed that VLCKD decreased BW, triglyceride and diastolic blood pressure while increased high-density lipoprotein (HDL-C) and low-density

lipoproteins (LDL-C) compared to LCD (Bueno *et al.*, 2013). Results of another meta-analysis study showed that LCD-C reduced triglycerides concentration and increased HDL cholesterol concentration (Meng *et al.*, 2017). A meta-analysis study showed that LCD can significantly decrease BMI and serum levels of total cholesterol and LDL-C in polycystic ovary syndrome (PCOS) women (Zhang *et al.*, 2019). Some reports have shown that LCD cannot have a protective effect (Foster *et al.*, 2003, Yancy Jr *et al.*, 2004). Results of a study have indicated that LCD can affect weight loss more than the conventional diet in the first six months, but the differences were not significant during a year in another study (Foster *et al.*, 2003). It was also shown that changes in LDL-C levels did not differ statistically with the LCD and LFD (Yancy Jr *et al.*, 2004).

The low percentage of carbohydrate intake from energy will be compensated with high fat and protein intake (Farabi and Hernandez, 2019, Frigolet *et al.*, 2011). Protein consists of amino acids. Amino acids are necessary for preserving lean body mass and muscle growth (Coker *et al.*, 2012, Simonson *et al.*, 2020), also intake of protein is positively associated with lean mass (Houston *et al.*, 2008). Ebbeling *et al.* reported that to decrease an individual's BMR (basal metabolic rate) LCD is effective compared to LFD. LCD may result in the modulation of resting energy expenditure (Ebbeling *et al.*, 2012). LCD decreases supply of glucose to the liver, muscles and brain, thereby inactivating gluconeogenesis. The main source of gluconeogenesis is glycerol (Brouns, 2018). It is caused by triglycerides breakdown (Chourpiliadis and Mohiuddin, 2020). The Decomposition of triglycerides is associated with reduced FM. The current study results showed that LCD significantly reduced body FM percentage; however, no significant difference in body FM changes was observed. In fact, LCD can change the body composition.

This study had several strengths. RCTs with less bias were included in comparison to observational studies. A large number of included studies lets us assess publication bias. This meta-analysis due to

the large sample size could recognize statistically significant mean differences in outcomes. Further subgroup analyses based on the duration of follow-up, gender, and health status would provide beneficial insights into these diets.

This study had some limitations. Given that one of the inclusion criteria was English language publications, some studies might have been missed. Most studies did not prepare food for the duration of the trial, which may reduce adherence to diet protocols. Some studies did not also report any data about participant's physical activity. Crossover trials in this study only reported overall results for low-carb and low-fat arm; so, these measures were used in the present analysis. Furthermore, diets did not account for the quality of the food consumed. The type of intake carbohydrates was not clear. Refine cereals or whole grains have different effects on the risk of disease. Other limitations of the study were heterogeneity during follow-up and the amount of carbohydrates in both the interventional and comparative diets.

Conclusion

Dietary intake is the main cause of abnormal anthropometric characteristics, and diet changes are suggested to improve this condition. Generally, LCD compared to LFD had an effect in losing BMI, weight and body fat percentage, but did not affect body FM and WC. In the future, more interventions with a specific carbohydrate dose are needed to reach a definitive conclusion.

Acknowledgment

Thanks are owed to the Research Committee of Kermanshah University of Medical Sciences.

Funding information

The present study with ID 3011110 was supported by the Nutrition Research Committee of School of Nutrition and Food Science Kermanshah University of Medical Sciences.

Conflict of interest

The authors declare that they have no conflict of interest.

Authors' contributions

Darand M designed the research; Darand M and Alizadeh S conducted the research; Hassanizadeh S analyzed the data; Darand M, Talebi S and Darand Z extracted the data, Darabi Z, Yaghoubi F, Bagherniya M and Azamian Y wrote the paper, Abdollahzad H finalized the paper. All authors read and approved the final manuscript.

References

- Abarca-Gómez L, et al.** 2017. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128· 9 million children, adolescents, and adults. *The Lancet*. **390 (10113)**: 2627-2642.
- Astrup A, Larsen TM & Harper A** 2004. Atkins and other low-carbohydrate diets: hoax or an effective tool for weight loss? *The Lancet*. **364 (9437)**: 897-899.
- Borenstein M, Hedges LV, Higgins JP & Rothstein HR** 2011. Introduction to meta-analysis. John Wiley & Sons.
- Brinkworth GD, Noakes M, Clifton PM & Bird AR** 2009. Comparative effects of very low-carbohydrate, high-fat and high-carbohydrate, low-fat weight-loss diets on bowel habit and faecal short-chain fatty acids and bacterial populations. *British journal of nutrition*. **101 (10)**: 1493-1502.
- Brouns F** 2018. Overweight and diabetes prevention: is a low-carbohydrate-high-fat diet recommendable? *European journal of nutrition*. **57 (4)**: 1301-1312.
- Buchholz AC & Schoeller DA** 2004. Is a calorie a calorie? *American journal of clinical nutrition*. **79 (5)**: 899S-906S.
- Bueno NB, de Melo ISV, de Oliveira SL & da Rocha Ataíde T** 2013. Very-low-carbohydrate ketogenic diet v. low-fat diet for long-term weight loss: a meta-analysis of randomised controlled trials. *British journal of nutrition*. **110 (7)**: 1178-1187.
- Burgess B, Raynor HA & Tepper BJ** 2017. PROP Nontaster Women Lose More Weight Following a Low-Carbohydrate Versus a Low-

- Fat Diet in a Randomized Controlled Trial. *Obesity (Silver Spring, Md.)*. **25 (10)**: 1682-1690.
- Chawla S, Tassarolo Silva F, Amaral Medeiros S, Mekary RA & Radenkovic D** 2020. The Effect of Low-Fat and Low-Carbohydrate Diets on Weight Loss and Lipid Levels: A Systematic Review and Meta-Analysis. *Nutrients*. **12 (12)**: 3774.
- Chourpiliadis C & Mohiuddin SS** 2020. Biochemistry, gluconeogenesis. *StatPearls* <https://www.ncbi.nlm.nih.gov/books/NBK544346/>
- Coker RH, Miller S, Schutzler S, Deutz N & Wolfe RR** 2012. Whey protein and essential amino acids promote the reduction of adipose tissue and increased muscle protein synthesis during caloric restriction-induced weight loss in elderly, obese individuals. *Nutrition journal*. **11 (1)**: 105.
- de Luis DA, et al.** 2007. Effects of a low-fat versus a low-carbohydrate diet on adipocytokines in obese adults. *Hormone research*. **67 (6)**: 296-300.
- Diabetes U** 2017. Position statement: low-carb diets for people with diabetes. *Diabetes UK: London, UK*.
- Duval S & Tweedie R** 2000. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*. **56 (2)**: 455-463.
- Ebbeling CB, et al.** 2012. Effects of dietary composition on energy expenditure during weight-loss maintenance. *Journal of the American medical association*. **307 (24)**: 2627-2634.
- Elmadfa I & Meyer AL** 2008. Body composition, changing physiological functions and nutrient requirements of the elderly. *Annals of nutrition and metabolism*. **52 (Suppl. 1)**: 2-5.
- Farabi SS & Hernandez TL** 2019. Low-carbohydrate diets for gestational diabetes. *Nutrients*. **11 (8)**: 1737.
- Foster GD, et al.** 2003. A randomized trial of a low-carbohydrate diet for obesity. *New England Journal of Medicine*. **348 (21)**: 2082-2090.
- Frigolet M-E, Barragán V-ER & González MT** 2011. Low-carbohydrate diets: a matter of love or hate. *Annals of nutrition and metabolism*. **58 (4)**: 320-334.
- Frisch S, et al.** 2009. A randomized controlled trial on the efficacy of carbohydrate-reduced or fat-reduced diets in patients attending a telemedically guided weight loss program. *Cardiovascular diabetology*. **8**: 36.
- Furtado JM, et al.** 2018. Anthropometric features as predictors of atherogenic dyslipidemia and cardiovascular risk in a large population of school-aged children. *PloS one*. **13 (6)**: e0197922.
- Gardner CD, et al.** 2007. Comparison of the Atkins, Zone, Ornish, and LEARN diets for change in weight and related risk factors among overweight premenopausal women: the A TO Z Weight Loss Study: a randomized trial. *Journal of American medical association*. **297 (9)**: 969-977.
- Gardner CD, et al.** 2018. Effect of low-fat VS low-carbohydrate diet on 12-month weight loss in overweight adults and the association with genotype pattern or insulin secretion the DIETFITS randomized clinical trial. *Journal of the American medical association*. **319 (7)**: 667-679.
- Hadaegh F, Shafiee G & Azizi F** 2009. Anthropometric predictors of incident type 2 diabetes mellitus in Iranian women. *Annals of Saudi medicine*. **29 (3)**: 194-200.
- Harcombe Z, et al.** 2015. Evidence from randomised controlled trials did not support the introduction of dietary fat guidelines in 1977 and 1983: a systematic review and meta-analysis. *Open heart*. **2 (1)**: e000196.
- Hashimoto Y, et al.** 2016. Impact of low-carbohydrate diet on body composition: meta-analysis of randomized controlled studies. *Obesity reviews*. **17 (6)**: 499-509.
- Haufe S, et al.** 2011. Randomized comparison of reduced fat and reduced carbohydrate hypocaloric diets on intrahepatic fat in overweight and obese human subjects. *Hepatology (Baltimore, Md.)*. **53 (5)**: 1504-1514.

- Higgins J** 2011. Cochrane handbook for systematic reviews of interventions. Version 5.1.0 [updated March 2011]. The Cochrane Collaboration.
- Higgins JP, et al.** 2019. Cochrane handbook for systematic reviews of interventions. John Wiley & Sons.
- Higgins JP & Thompson SG** 2002. Quantifying heterogeneity in a meta-analysis. *Statistics in medicine*. **21** (11): 1539-1558.
- Hite AH, Berkowitz VG & Berkowitz K** 2011. Low-carbohydrate diet review: shifting the paradigm. *Nutrition in clinical practice*. **26** (3): 300-308.
- Houston DK, et al.** 2008. Dietary protein intake is associated with lean mass change in older, community-dwelling adults: the Health, Aging, and Body Composition (Health ABC) Study. *American journal of clinical nutrition*. **87** (1): 150-155.
- Jang HH, et al.** 2017. Agrimonia pilosa Ledeb. aqueous extract improves impaired glucose tolerance in high-fat diet-fed rats by decreasing the inflammatory response. *BMC complementary and alternative medicine*. **17** (1).
- Janssen F, Bardoutsos A & Vidra N** 2020. Obesity Prevalence in the Long-Term Future in 18 European Countries and in the USA. *Obesity facts*. **13** (5): 514-527.
- Keys A, et al.** 1986. The diet and 15-year death rate in the seven countries study. *American journal of epidemiology*. **124** (6): 903-915.
- Khader Y, Batieha A, Jaddou H, El-Khateeb M & Ajlouni K** 2019. The performance of anthropometric measures to predict diabetes mellitus and hypertension among adults in Jordan. *BMC public health*. **19** (1): 1-9.
- Lichtenstein AH & Van Horn L** 1998. Very low fat diets. *Circulation*. **98** (9): 935-939.
- Lodi A, Zarantonello L, Bisiacchi PS, Cenci L & Paoli A** 2020. Ketonemia and Glycemia Affect Appetite Levels and Executive Functions in Overweight Females During Two Ketogenic Diets. *Obesity*. **28** (10): 1868-1877.
- Lorzadeh E, Ramezani-Jolfaie N, Mohammadi M, Khoshtakht Y & Salehi-Abargouei A** 2019. The effect of hesperidin supplementation on inflammatory markers in human adults: a systematic review and meta-analysis of randomized controlled clinical trials. *Chemico-biological interactions*. **307**: 8-15.
- Mansoor N, Vinknes KJ, Veierød MB & Retterstøl K** 2016. Effects of low-carbohydrate diets v. low-fat diets on body weight and cardiovascular risk factors: a meta-analysis of randomised controlled trials. *British journal of nutrition*. **115** (3): 466-479.
- Meng Y, et al.** 2017. Efficacy of low carbohydrate diet for type 2 diabetes mellitus management: a systematic review and meta-analysis of randomized controlled trials. *Diabetes research and clinical practice*. **131**: 124-131.
- Moher D, et al.** 2015. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic reviews*. **4** (1): 1.
- Noakes M, Keogh JB, Foster PR & Clifton PM** 2005. Effect of an energy-restricted, high-protein, low-fat diet relative to a conventional high-carbohydrate, low-fat diet on weight loss, body composition, nutritional status, and markers of cardiovascular health in obese women. *American journal of clinical nutrition*. **81** (6): 1298-1306.
- Nordmann AJ, et al.** 2006. Effects of low-carbohydrate vs low-fat diets on weight loss and cardiovascular risk factors: a meta-analysis of randomized controlled trials. *Archives of internal medicine*. **166** (3): 285-293.
- Perna S, et al.** 2019. Effectiveness of a hypocaloric and low-carbohydrate diet on visceral adipose tissue and glycemic control in overweight and obese patients with type 2 diabetes. *Bahrain medical bulletin*. **41** (3): 159-164.
- Petrisko M, et al.** 2020. Biochemical, Anthropometric, and Physiological Responses to Carbohydrate-Restricted Diets Versus a Low-Fat Diet in Obese Adults: A Randomized Crossover Trial. *Journal of medicinal food*. **23** (3): 206-214.
- Phillips SA, et al.** 2008. Benefit of low-fat over low-carbohydrate diet on endothelial health in

- obesity. *Hypertension (Dallas, Tex. : 1979)*. **51** (2): 376-382.
- Rajaie S, Azadbakht L, Saneei P, Khazaei M & Esmailzadeh A** 2013. Comparative effects of carbohydrate versus fat restriction on serum levels of adipocytokines, markers of inflammation, and endothelial function among women with the metabolic syndrome: a randomized cross-over clinical trial. *Annals of nutrition and metabolism*. **63** (1-2): 159-167.
- Rock CL, et al.** 2016. Effects of diet composition on weight loss, metabolic factors and biomarkers in a 1-year weight loss intervention in obese women examined by baseline insulin resistance status. *Metabolism: clinical and experimental*. **65** (11): 1605-1613.
- Sackner-Bernstein J, Kanter D & Kaul S** 2015. Dietary intervention for overweight and obese adults: comparison of low-carbohydrate and low-fat diets. A meta-analysis. *PLoS one*. **10** (10): e0139817.
- Schwingshackl L & Hoffmann G** 2013. Long-term effects of low-fat diets either low or high in protein on cardiovascular and metabolic risk factors: a systematic review and meta-analysis. *Nutrition journal*. **12** (1): 48.
- Sharman MJ & Volek JS** 2004. Weight loss leads to reductions in inflammatory biomarkers after a very-low-carbohydrate diet and a low-fat diet in overweight men. *Clinical science*. **107** (4): 365-369.
- Simonson M, Boirie Y & Guillet C** 2020. Protein, amino acids and obesity treatment. *Reviews in endocrine and metabolic disorders*. **21** (3): 341-353.
- Tobias DK, et al.** 2015. Effect of low-fat diet interventions versus other diet interventions on long-term weight change in adults: a systematic review and meta-analysis. *Lancet Diabetes & endocrinology*. **3** (12): 968-979.
- van Zuuren EJ, Fedorowicz Z, Kuijpers T & Pijl H** 2018. Effects of low-carbohydrate- compared with low-fat-diet interventions on metabolic control in people with type 2 diabetes: a systematic review including GRADE assessments. *American journal of clinical nutrition*. **108** (2): 300-331.
- Vander Wal JS, Mccburney MI, Moellering N, Marth J & Dhurandhar NV** 2007. Moderate-carbohydrate low-fat versus low-carbohydrate high-fat meal replacements for weight loss. *International journal of food sciences and nutrition*. **58** (4): 321-329.
- Wadden TA, Butryn ML & Wilson C** 2007. Lifestyle modification for the management of obesity. *Gastroenterology*. **132** (6): 2226-2238.
- Wadden TA, Webb VL, Moran CH & Bailer BA** 2012. Lifestyle modification for obesity: new developments in diet, physical activity, and behavior therapy. *Circulation*. **125** (9): 1157-1170.
- Wycherley TP, et al.** 2010. Long-term effects of weight loss with a very low carbohydrate and low fat diet on vascular function in overweight and obese patients: Original Article. *Journal of internal medicine*. **267** (5): 452-461.
- Yancy Jr WS, Olsen MK, Guyton JR, Bakst RP & Westman EC** 2004. A low-carbohydrate, ketogenic diet versus a low-fat diet to treat obesity and hyperlipidemia: a randomized, controlled trial. *Annals of internal medicine*. **140** (10): 769-777.
- Zhang L, et al.** 2020. Prevalence of overweight and obesity in China: results from a cross-sectional study of 441 thousand adults, 2012–2015. *Obesity research & clinical practice*. **14** (2): 119-126.
- Zhang X, Zheng Y, Guo Y & Lai Z** 2019. The effect of low carbohydrate diet on polycystic ovary syndrome: a meta-analysis of randomized controlled trials. *International journal of endocrinology*. **2019**.