



# Journal of Nutrition and Food Security

Shahid Sadoughi University of Medical Sciences  
School of Public Health  
Department of Nutrition  
Nutrition & Food Security Research Center



eISSN: 2476-7425

pISSN: 2476-7417

JNFS 2017; 2(4): 262-264

Website: jnfs.ssu.ac.ir

## *Metabolic Syndrome: An Important Health Issue in Iran*

Mohammad Mohammadi; MSc<sup>1,2</sup> & Amin Salehi-Abargouei; PhD<sup>1,2\*</sup>

<sup>1</sup> Nutrition and Food Security Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

<sup>2</sup> Department of Nutrition, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

### ARTICLE INFO

#### EDITORIAL ARTICLE

#### Article history:

Received: 17 Nov 2016

Revised: 21 Dec 2016

Accepted: 21 Jan 2017

#### Corresponding author:

Department of Nutrition, Faculty of Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

abargouei@ssu.ac.ir

Postal code: 8916877391

Tel: +98-35-31492229

Metabolic syndrome (MetS) is a major metabolic disorder worldwide which is characterized by having three or more of the five metabolic abnormalities. These metabolic abnormalities include abdominal obesity (waist circumference  $\geq 90$  cm for men and  $\geq 80$  cm for women), high triglyceride levels (TG) ( $\geq 150$  mg/dL), decreased high density lipoprotein cholesterol (HDL-c) ( $< 40$  mg/dL in men and  $< 50$  mg/dL in women), increased blood pressure (BP) (systolic  $\geq 130$  mmHg and/or diastolic  $\geq 85$  mmHg), and elevated fasting blood glucose (FBG) ( $\geq 100$  mg/dL) (Mohammadi *et al.*, 2016). These conditions induce insulin resistance, inflammation, and chronic stress resulting in excessive flux of fatty acids which can lead to increased risk for type 2 diabetes and cardiovascular diseases (Brunner *et al.*, 2002, Eckel *et al.*, 2005, Yaffe *et al.*, 2004). Approximately 20% of the patients with MetS who do not have diabetes remain at high risk of incident cardiovascular events (Wong *et al.*, 2003a, Wong

*et al.*, 2003b). The population-attributable fraction for all-cause mortality from the MetS was 6–7% (Ford, 2005).

In 2012, the total costs of diagnosed diabetes in the USA were estimated to be around 245 billion dollars; US \$176 billion of which were direct medical costs (Association, 2013). Furthermore, the American Heart Association reported that the one-year cost of cardiovascular diseases (CVD) is US \$457.4 billion in the USA (Thom *et al.*, 2006). In a survey conducted among the Iranian population, the costs of essential diagnostic and therapeutic services of CVD patients in the Central Hospital of National Iranian Oil Corporation were nearly 10.94 billion Rials (Fakhzadeh *et al.*, 2002). Therefore, regarding the heavy economic burden of MetS and other associated diseases on the health care system, it is necessary to pay special attention to its prevention, control, and treatment.

The prevalence of MetS was reported to be around 24% in the United States, 12% in Europe, and 10–40% in most Asian countries (Ryan *et al.*,

**This paper should be cited as: Mohammadi M, Salehi-Abargouei A. Metabolic Syndrome: an important health issue in Iran. Journal of Nutrition and Food Security (JNFS), 2017; 2 (4): 262-264.**

2008). In Iran, the rate of MetS was estimated to be approximately 36.9%, 34.6%, and 41.5% according to the Adult Treatment Panel III (ATP III) criteria, the International Diabetes Federation (IDF) criteria, and the Joint Interim Societies (JIS) criteria, respectively (Amirkalali *et al.*, 2015). It was also reported that more than 32% of population of Yazd city have MetS (Sadrbafoghi *et al.*, 2006). Another survey among female teachers in Yazd showed that the prevalence of MetS was high even in the educated people of this area (39.11% based on ATP III definition) (Shahvazi *et al.*, 2016).

The MetS complications and comorbid diseases can be prevented by appropriate control of body weight, lipid profile, and blood pressure (Malik *et al.*, 2004). Indeed, it was indicated that all components of MetS may be related to obesity. Therefore, weight loss and physical activity might be the most important strategies to improve the MetS risk factors (Olefsky *et al.*, 1974).

## References

- Amirkalali B, et al.** 2015. Prevalence of Metabolic Syndrome and Its Components in the Iranian Adult Population: A Systematic Review and Meta-Analysis. *Iranian red crescent medical journal.* **17 (12):** e24723.
- Association AD** 2013. Economic costs of diabetes in the US in 2012. *Diabetes Care* 2013; **36:** 1033–1046. *Diabetes care.* **36 (6):** 1797.
- Azadbakht L, Mirmiran P, Esmailzadeh A, Azizi T & Azizi F** 2005. Beneficial effects of a Dietary Approaches to Stop Hypertension eating plan on features of the metabolic syndrome. *Diabetes care.* **28 (12):** 2823-2831.
- Babio N, Bulló M & Salas-Salvadó J** 2009. Mediterranean diet and metabolic syndrome: the evidence. *Public health nutrition.* **12 (9A):** 1607-1617.
- Brunner EJ, et al.** 2002. Adrenocortical, autonomic, and inflammatory causes of the metabolic syndrome: nested case-control study. *Circulation.* **106 (21):** 2659-2665.
- Eckel RH, Grundy SM & Zimmet PZ** 2005. The metabolic syndrome. *Lancet.* **365 (9468):** 1415-1428.
- Fakhzadeh H, Mohaghegh M, Poor-Ebrahim R & Akhlaghi M** 2002. The economic cost of coronary heart disease in iran oil industry personnel in 1999. *Journal of medical council of I.R.I.* **19 (4):** 287-292.
- Feldeisen SE & Tucker KL** 2007. Nutritional strategies in the prevention and treatment of metabolic syndrome. *Applied physiology, nutrition, and metabolism.* **32 (1):** 46-60.
- Ford ES** 2005. Risks for all-cause mortality, cardiovascular disease, and diabetes associated with the metabolic syndrome a summary of the evidence. *Diabetes care.* **28 (7):** 1769-1778.
- Malik S, et al.** 2004. Impact of the metabolic syndrome on mortality from coronary heart disease, cardiovascular disease, and all causes in United States adults. *Circulation.* **110 (10):** 1245-1250.

Consumption of healthy diets containing more vegetables, fruits, whole grains, low-fat dairy products, and replacing some dietary carbohydrates with monounsaturated fat (e.g. olive oil) may also be beneficial for most patients with MetS (Feldeisen and Tucker, 2007, Prasad *et al.*, 2012). Healthy dietary patterns such as Dietary Approaches to Stop Hypertension (DASH) and Mediterranean diet can decrease most metabolic risk factors in both men and women. Indeed, these diets were inversely associated with Body mass index which can be useful for improvement of hypertension, atherogenic dyslipidemia, or inflammatory reactions associated with MetS (Azadbakht *et al.*, 2005, Babio *et al.*, 2009).

Unfortunately, the health care system in Iran is more focused on treatment of diseases than implementation of preventive strategies. Planning national programs which benefit from intersectoral collaborations for determining and controlling the MetS risk factors seems to be urgent in Iran.

- Mohammadi M, Jolfaie NR, Alipour R & Zarrati M** 2016. Is Metabolic Syndrome Considered to Be a Risk Factor for Gastroesophageal Reflux Disease (Non-Erosive or Erosive Esophagitis)? A Systematic Review of the Evidence. *Iranian red crescent medical journal*. **18 (11)**: e30363.
- Olefsky J, Reaven GM & Farquhar JW** 1974. Effects of weight reduction on obesity studies of lipid and carbohydrate metabolism in normal and hyperlipoproteinemic subjects. *Journal of clinical Investigation*. **53 (1)**: 64.
- Prasad H, Ryan DA, Celzo MF & Stapleton D** 2012. Metabolic syndrome: definition and therapeutic implications. *Postgraduate medicine*. **124 (1)**: 21-30.
- Ryan AM, et al.** 2008. Barrett esophagus: prevalence of central adiposity, metabolic syndrome, and a proinflammatory state. *Annals of surgery*. **247 (6)**: 909-915.
- Sadrbafoghi S, et al.** 2006. Prevalence and criteria of metabolic syndrome in an urban population: Yazd Healthy Heart Project. *Tehran university medical journal*. **64 (10)**: 90-96.
- Shahvazi S, Nadjarzadeh A, Mehri Z & Salehi-Abargouei A** 2016. Prevalence of Metabolic Syndrome in Adult Females: Comparison between Iranian National Definition and Currently Used International Criteria. *Journal of nutrition and food security* **1(1)**: 49-62.
- Thom T, et al.** 2006. Heart disease and stroke statistics--2006 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation*. **113 (6)**: e85-151.
- Wong ND, et al.** 2003a. Preventing coronary events by optimal control of blood pressure and lipids in patients with the metabolic syndrome. *The American journal of cardiology*. **91 (12)**: 1421-1426.
- Wong ND, et al.** 2003b. The metabolic syndrome, diabetes, and subclinical atherosclerosis assessed by coronary calcium. *Journal of the American College of Cardiology*. **41 (9)**: 1547-1553.
- Yaffe K, et al.** 2004. The metabolic syndrome, inflammation, and risk of cognitive decline. *Journal of the American medical association*. **292 (18)**: 2237-2242.