



Dietary Fats, Minerals and Semen Quality and Quantity among Men Attending a Fertility Clinic: A Cross-Sectional Study in Iran

Mehran Rahimlou; PhD¹, Sara Sohaei; MSc², Ammar Salehi-Sahlabadi; PhD³ & Mehran Nouri; PhD *^{4,5,6}

¹ Department of Nutrition, School of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran; ² Department of Clinical Nutrition, School of Nutrition and Food Science, Isfahan University of Medical Sciences, Isfahan, Iran; ³ Student Research Committee, Department of Clinical Nutrition and Dietetics, School of Nutrition and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran; ⁴ Students Research Committee, Shiraz University of Medical Sciences, Shiraz, Iran; ⁵ Department of Community Nutrition, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran; ⁶ Health Policy Research Center, Institute of Health, Shiraz University of Medical Sciences, Shiraz, Iran.

ARTICLE INFO

ORIGINAL ARTICLE

Article history:

Received: 7 Sep 2021

Revised: 21 Feb 2022

Accepted: 3 Mar 2022

*Corresponding author:

mehran_nouri71@yahoo.com
Department of Community Nutrition, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran.

Postal code: 71348-14336

Tel: +98 7137258099

ABSTRACT

Background: At least 50% of infertile couple's problems are related to male factor infertility and in many patients; nutritional problems have been related to decreased sperm quality. Thus, the present study aims to examine the association of dietary fat and mineral intake with semen quantity and quality. **Methods:** This cross-sectional study was conducted on 175 infertile men in Isfahan Infertility and Fertility Center, Iran. Dietary intake of all patients was evaluated by validated Food Frequency Questionnaire (FFQ) and semen analysis was performed by standard protocol. Linear and quartile regression were used to determine the association of dietary fat and mineral intake with semen related factors. **Results:** There was a positive association between the sperm total count with iron (Fe) ($P_{trend} = 0.03$), zinc (Zn) ($P_{trend} = 0.02$), and selenium (Se) ($P_{trend} = 0.001$). Men in the highest quartile of Zn and alpha-linolenic acid (ALA) had higher sperm density ($P < 0.05$). Levels of Zn and docosahexaenoic acid (DHA) in participant's diet were also negatively related to DNA damage ($P_{trend} = 0.01$ for Zn and $P_{trend} = 0.04$ for DHA). Higher intake of saturated fatty acid (SFA) was associated with lower sperm density ($P_{trend} = 0.02$) and higher intake of monounsaturated fatty acids (MUFAs) was associated with higher sperm totality ($P_{trend} = 0.02$). **Conclusion:** In this preliminary cross-sectional study, diets containing higher amounts of Fe, Zn, Se, MUFA, and DHA and lower amounts of SFA were associated with more favourable semen quality parameters.

Keywords: Male infertility; Semen quality, Minerals, Fats

Introduction

Over the past years, one of the concerns of health professionals has been increasing the incidence of infertility in both genders (Irvine *et al.*, 1996, Zou *et al.*, 2011). More than 70 million

couples suffer from infertility worldwide. More than 50% of infertility is associated with male disorders (Mehra *et al.*, 2018). Contradictory results have been reported regarding the decrease

This paper should be cited as: Rahimlou M, Sohaei S, Salehi-Sahlabadi A, Nouri M. Dietary Fats, Minerals and Semen Quality and Quantity among Men Attending a Fertility Clinic: A Cross-Sectional Study in Iran. Journal of Nutrition and Food Security (JNFS), 2022; 7(4): 452-463.

in the quality and quantity of sperm in young men (Auger *et al.*, 1995, Rolland *et al.*, 2012). Researchers have suggested that reducing sperm quality is more affected by environmental factors. Some factors, such as radiation, smoking, varicocele, infection, and urinary tract infection contribute to the etiology of male infertility. Nevertheless, regional differences also affect the results of various studies (Jorgensen *et al.*, 2001, Lipshultz, 1996, Paulsen *et al.*, 1996, Wong *et al.*, 2000), so that, infertility rates are higher in developed and industrialized countries (Danielewicz *et al.*, 2018). One of the environmental factors affecting the quality of sperm is individuals' diets. Little is known of how diet may affect male reproductive potential. Emerging literature supports the hypothesis that specific nutritional factors can affect semen quality. Previous studies have shown that some food components, such as antioxidants, fruits, and vegetables can affect the quality of sperm (Afeiche *et al.*, 2013, Eslamian *et al.*, 2012). Other factors, such as the high consumption of alcoholic beverages, obesity, coffee, processed meat, and fast foods can also reduce sperm quality (Salas-Huetos *et al.*, 2017, Sermondade *et al.*, 2013).

Dietary fats are one of the essential dietary ingredients. Cholesterol as a type of dietary fat plays an important role in the structure of mammalian membranes (Diaz-Fontdevila and Bustos-Obregon, 1993). The sperm membrane is affected by the type of dietary fat (Saez Lancellotti *et al.*, 2010). Overconsumption of dietary saturated and Trans fatty acids by increasing inflammation in the body may affect sperm quality (Dadkhah *et al.*, 2017). On the other hand, previous studies have shown that there is an important association between the intake of certain dietary minerals (Ammar *et al.*, 2019, Fallah *et al.*, 2018, Moslemi and Tavanbakhsh, 2011), zinc (Zn), iron (Fe), and semen quality.

Although some previous studies have evaluated the association between the dietary components and quality of sperm, the results of existing studies are contradictory. Therefore, this study aims to investigate the relation between dietary fats dietary

fats and minerals with factors related to semen quantity and quality.

Materials and Methods

Study design: The present study was conducted using a case-control design on 175 infertile participants referred to Isfahan Infertility and Fertility Center, Iran, in 2018. The study participants were infertile men. The inclusion criteria consisted of adults aged 20-45 years, sperm counts lower than 20 million, motility lower than 60%, damaged morphology greater than 65%. The exclusion criteria included frequent consumption of omega 3, vitamin E and other antioxidant supplements, history of reproductive disorders or vasectomy, smoking, urinary tract infection, testicular surgery, testicular tumor, anatomical abnormalities of the genital area (Complete information has been previously published (Rahimlou *et al.*, 2019)).

Measurements: A bioelectrical impedance analysis (Omron BF511; Omron Healthcare, Osaka, Japan) was employed to measure weight with minimum clothes and without shoes. Height was measured using a Seca stadiometer with the precision of 0.1 cm in a standing position next to a wall, without shoes, and with relaxed shoulders. Then body mass index (BMI) was computed by dividing the weight (kg) by the square of height (m²).

For evaluation of dietary intake, a validated detailed food frequency questionnaire (FFQ) was used (Esmailzadeh *et al.*, 2007). This questionnaire contains 168 food items that are consumed more by the Iranian population. The participants were asked to report on average, the frequency of food intake variation from never or less than once/month to 12 times or more per day, during the previous year. Eventually, the amount of consumption of each item of food was computed and converted to grams per day using household measures (Ghaffarpour *et al.*, 1999) (Complete information has been previously published (Rahimlou *et al.*, 2019)).

Semen analysis: Semen samples were collected from all of the participants by masturbation. Of all

the participants, two samples of sperm were taken and if the difference between the samples was more than 20%, the third sample was taken. The samples were held at 37 °C to liquefy. After liquefaction, the samples were analyzed according to the World Health Organization guideline (Künzle *et al.*, 2003). Semen counts and motility percentage were evaluated using computer-aided semen analysis (CASA) (Hamilton-Thorne Version 10HTM-IVOS) using 2 Chamber Lejawa slides (Leja, The Netherlands). Morphology was established according to the Kruger parameters. Finally, sperm DNA damage was assessed by terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) and the sperm chromatin structure assay (SCSA).

Ethical considerations: The study was approved by the Ethics Committee from Esfahan University of Medical Sciences with the code of IR.MUI.RESEARCH.REC.1397.267 and informed consent was obtained from all the participants. Prior to data collection, the purpose of the study was explained to the participants and their informed consent was recorded.

Data analysis: Sperm samples were taken from 175 infertile men. These men were divided into four groups according to quartile of intake of total major fat and minerals categories. Participants with the lowest intake of each fats or minerals were considered as the reference group. Linear regression was used to examine the association of fats or minerals intake with semen quality parameters. Analysis of covariance (ANCOVA) was used for calculating adjusted semen parameters for each nutrient quartile by relevant covariates. Multivariate ANCOVA models were created with continuous semen parameters as dependent variables, and fats and minerals covariates as independent variables. For statistical analyses, SPSS® 20 (IBM Corp, Armonk, NY) was used and P-value < 0.05 was considered significant.

Results

A total of 175 participants aged from 20 to 45 years were included in this cross-sectional study.

Baseline characteristics of the participants are summarized in **Table 1**. The participants were infertile men with mean age of 32.19 ± 2.34 year and BMI of 26.86 ± 1.60 kg/m². Thirty-seven percent of the participants were smokers, and five percent consumed alcohol. All of the participants performed semen analysis and semen analysis parameters, including the mean sperm concentration was 10.39 ± 32.5 mill/mL; seminal volume, 4.20 ± 1.89 ml, and total sperm count was 42.91 ± 30.91 million. Dietary intake of the participants is shown in the **Table 1**. The mean daily total intake was 2183.56 ± 202.81 kcal/day for total energy, 153.23 ± 11.85 g/day for carbohydrate, 88.21 ± 10.40 g/day for protein, and 75.34 ± 13.41 g/day for total fat. The mean caffeine intake was 65.45 ± 46.01 mg/day.

The correlation between sperm related parameters with dietary components is shown in **Table 2**. Based on the table, there was no significant association between dietary intake of carbohydrate, protein, total fat, and caffeine with none of the sperm related factors ($P > 0.05$).

The mean daily intake of dietary minerals was 1212.49 ± 432.45 mg/day for calcium, 16.5 ± 4.32 mg/day for Fe, 425.86 ± 114.28 mg/day for magnesium, 14.35 ± 4.41 for Zn, 1.93 ± 0.58 µg/day, and 101.52 ± 26.58 µg/day for selenium (Se). According to **Table 3**, after adjusting for covariates, the sperm total count had linear association with Fe ($P_{trend} = 0.03$), Zn ($P_{trends} = 0.02$) and Se intake ($P_{trend} = 0.001$). The participants in the higher quartile of Zn consumption had higher sperm density ($P_{trend} = 0.03$). Moreover, a linear association was found between Se intake with sperm total motility ($P_{trend} = 0.01$) and an inverse association between Zn intake with sperm DNA damage ($P_{trend} = 0.01$).

The mean daily intake of dietary fats was 338.19 ± 221.46 mg/day for total cholesterol, 28.17 ± 11.13 mg/day for saturated fatty acid (SFA), 28.38 ± 11.91 mg/day for monounsaturated fatty acid (MUFA), 16.75 ± 7.66 mg/day for polyunsaturated fatty acid (PUFA), 14.04 ± 6.99 mg/day for linoleic acid, 1.27 ± 0.67 mg/day for alpha-linolenic acid (ALA), 0.07 ± 0.5 mg/day for

ecosapentaenoic acid (EPA), and 0.25 ± 0.23 mg/day for docosahexaenoic acid (DHA).

Table 4 reveals that there was an inverse significant association between dietary intake of SFA and semen density ($P_{trend} = 0.02$). A significant linear association was also found between MUFA intake and sperm total motility ($P_{trend} = 0.02$), ALA with sperm density ($P_{trend} =$

0.04) and sperm total motility ($P_{trend} = 0.01$), and between dietary DHA with sperm density ($P_{trend} = 0.03$). Finally, the subjects in the higher quartile of dietary intake of DHA had lower percentage of DNA damage ($P_{trend} = 0.04$) and the participants in the higher quartile of PUFA had lower percentage of DNA fragmentation index (DFI) ($P_{trend} = 0.03$).

Table 1. Baseline characteristics of the study population.

Characteristics	Total (n = 57)
Age (year)	32.19 \pm 2.34 ^a
Height (cm)	178.17 \pm 4.12
Weight (kg)	85.28 \pm 5.51
Body mass index (kg/m ²)	26.86 \pm 1.60
MET (MET-h/week)	30.91 \pm 1.81
Dietary intake	
Energy (kcal/day)	2183.56 \pm 202.81
Carbohydrate (g/day)	153.23 \pm 11.85
Protein (g/day)	88.21 \pm 10.40
Fat (g/day)	75.34 \pm 13.41
Caffeine (mg/day)	65.45 \pm 46.01
Semen parameters	
Seminal volume (ml)	4.20 \pm 1.89
Sperm concentration (mill/ml)	10.39 \pm 32.50
Total sperm count (mill)	42.91 \pm 30.91
Motile sperm (PR+NP) (%)	25.65 \pm 16.76
Normal morphology (%)	2.31 \pm 1.32
DNA fragmentation index (%)	18.57 \pm 6.94

^a: Mean \pm SD

Table 2. Correlation between sperm related parameters with dietary components.

Sperm parameters	Energy (kcal/day)	Protein (g/day)	Carbohydrate (g/day)	Fat (g/day)	Caffeine (mg/day)
Total sperm count	0.08 (0.27)	0.9 (0.23)	0.15 (0.04)	-0.017 (0.82)	-0.07 (0.32)
Sperm density (10 ⁶ /ml)	0.012 (0.87)	-0.013 (0.86)	0.09 (0.23)	-0.07 (0.33)	-0.04 (0.16)
Total motility (%)	0.07 (0.34)	0.04 (0.6)	0.13 (0.07)	0.01 (0.87)	- 0.076 (0.55)
DNA damage (%)	-0.08 (0.26)	-0.08 (0.24)	-0.1 (0.17)	-0.04 (0.59)	-0.02 (0.17)
DNA fragmentation index (%)	-0.09 (0.22)	-0.08 (0.26)	-0.12 (0.1)	-0.04 (0.52)	-0.05 (0.48)

Table 3. Multivariate-adjusted model of dietary intake of minerals and semen parameters.

Variables	Sperm total count				Sperm density				Total motility				DNA damage				DNA Fragmentation Index			
	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b
Calcium																				
Q1																				
Q2	-0.77	6.61	0.9	0.69	1.55	1.27	0.23	0.24	1.70	3.57	0.62	0.81	-0.98	1.44	0.49	0.92	-0.76	1.48	0.6	0.95
Q3	8.13	6.57	0.21	0.55	2.1	1.28	0.1	0.18	4.79	3.59	0.18	0.57	0.96	1.45	0.5	0.13	0.89	1.49	0.54	0.17
Q4	7.34	6.54	0.26	0.7	0.68	1.29	0.59	0.69	2.61	3.61	0.46	0.89	-0.7	1.46	0.62	0.47	-0.75	1.48	0.63	0.64
P _{trend}		0.17				0.38				0.39				0.34				0.46		
Fe																				
Q1																				
Q2	3.69	6.41	0.56	0.69	-2.46	1.24	0.04	0.21	2.55	3.53	0.47	0.57	-2.29	1.44	0.62	0.66	-1.41	1.47	0.33	0.31
Q3	0.75	6.38	0.9	0.85	0.56	1.25	0.65	0.44	-0.93	3.51	0.78	0.7	-1.52	1.43	0.28	0.44	-1.95	1.46	0.17	0.27
Q4	15	6.41	0.01	0.18	1.26	1.26	0.31	0.2	6.58	3.53	0.06	0.41	-0.7	1.44	0.11	0.42	-2.74	1.47	0.06	0.26
P _{trend}		0.03				0.24				0.08				0.15				0.22		
Zn																				
Q1																				
Q2	2.45	4.28	0.32	0.42	0.76	2.75	0.56	0.63	1.19	2.16	0.20	0.36	-0.28	1.34	0.84	0.58	0.19	1.47	0.59	0.45
Q3	3.17	4.28	0.09	0.12	1.32	2.73	0.25	0.27	1.65	2.17	0.08	0.12	-0.94	1.34	0.32	0.22	-1.26	1.47	0.19	0.08
Q4	6.47	4.29	0.03	0.04	3.63	2.76	0.04	0.05	1.88	2.19	0.12	0.24	-1.68	1.35	0.09	0.04	-1.82	1.48	0.11	0.032
P _{trend}		0.02				0.03				0.16				0.01				0.04		
Cu																				
Q1																				
Q2	-1.97	1.26	0.11	0.3	7.69	3.52	0.02	0.007	-4.96	7.08	0.48	0.11	-1.18	1.66	0.46	0.3	-1.80	1.46	0.21	0.43
Q3	0.67	1.25	0.59	0.41	1.69	3.50	0.62	0.16	0.26	7.84	0.97	0.5	-1.87	1.77	0.28	0.12	-2.80	1.47	0.05	0.21
Q4	1.06	1.23	0.4	0.21	0.31	3.51	0.92	0.27	11.74	9.69	0.22	0.4	-0.87	2.19	0.69	0.19	-1.99	1.48	0.17	0.79
P _{trend}		0.18				0.73				0.44				0.16				0.58		
Se																				
Q1																				
Q2	1.66	6.51	0.37	0.77	-0.45	1.29	0.72	0.53	0.81	3.42	0.00	0.006	-0.39	1.44	0.23	0.08	-0.7	1.47	0.59	0.22
Q3	5.83	6.44	0.04	0.08	0.12	1.27	0.92	0.73	2.42	3.38	0.08	0.12	-1.63	1.43	0.24	0.78	-1.66	1.48	0.25	0.75
Q4	10.33	6.55	<0.001	0.001	0.27	1.28	0.83	0.94	5.42	3.44	0.03	0.048	-1.39	1.43	0.78	0.51	-1.85	1.45	0.97	0.47
P _{trend}		0.001				0.77				0.01				0.35				0.44		

P^a: crude p- value, P^b: adjusted p-value, tests for linear trend were performed using the median value for each quartile. Multivariate model adjusted for BMI, energy intake and physical activity.

Table 4. Multivariate-adjusted model of dietary intake of fats and semen parameters

Median for each quartile	Sperm total count				Sperm density				Total motility				DNA damage				DNA Fragmentation Index			
	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b
Cholesterol																				
Q1																				
Q2	-3.76	7.08	0.59	0.72	-0.63	1.27	0.61	0.76	-2.77	3.50	0.42	0.61	-0.71	1.58	0.65	0.48	-2.17	1.47	0.14	0.2
Q3	-12.59	7.63	0.1	0.32	-2.12	1.26	0.94	0.37	-9.63	3.48	0.006	0.03	0.69	1.70	0.68	0.73	-1.22	1.46	0.4	0.56
Q4	-11.52	9.00	0.201	0.46	-1.44	1.27	0.25	0.16	-1.27	3.50	0.71	0.48	-2.68	1.44	0.06	0.24	0.31	1.47	0.83	0.87
P _{trend}	0.38				0.28				0.4				0.53				0.16			
Saturated fatty acid (SFA)																				
Q1	-																			
Q2	-0.62	6.57	0.43	0.51	-0.6	1.28	0.63	0.24	-5.12	3.56	0.15	0.06	0.01	1.45	0.99	0.41	0.16	1.47	0.91	0.37
Q3	0.97	6.57	0.88	0.05	-1.15	1.25	0.12	0.21	-2.49	3.56	0.48	0.16	-1.15	1.45	0.42	0.68	-2.02	1.47	0.17	0.99
Q4	-5.11	6.53	0.43	0.69	-1.90	1.17	0.03	0.04	-1.48	3.54	0.67	0.27	0.15	1.44	0.91	0.2	0.29	1.46	0.84	0.72
P _{trend}	0.13				0.02				0.93				0.91				0.85			
Mono unsaturated fatty acid (MUFA)																				
Q1																				
Q2	2.57	7.16	0.72	0.9	-0.76	1.32	0.56	0.4	1.84	3.70	0.3	0.06	-1.68	1.50	0.26	0.72	-0.95	1.54	0.53	0.96
Q3	-2.13	7.13	0.76	0.89	-2.16	1.38	0.11	0.25	4.45	3.87	0.25	0.02	-1.26	1.57	0.41	0.85	-0.9	1.61	0.57	0.68
Q4	2.5	6.82	0.7	0.27	-1.24	1.39	0.37	0.82	5.28	3.89	0.29	0.04	-0.73	1.57	0.64	0.59	-0.5	1.61	0.75	0.51
P _{trend}	0.63				0.69				0.02				0.41				0.29			
Poly unsaturated fatty acid (PUFA)																				
Q1																				
Q2	1.32	6.43	0.01	0.05	-1.39	-0.5	1.27	0.69	4.16	3.54	0.24	0.32	-2.27	1.43	0.11	0.08	0.47	1.46	0.09	0.06
Q3	5.51	6.50	0.001	0.001	-0.54	1.15	1.29	0.9	-0.57	3.56	0.87	0.85	-1.59	1.44	0.27	0.11	-1.62	1.47	0.27	0.09
Q4	11.43	6.43	0.001	0.001	0.2	1.12	1.27	0.37	3.12	3.54	0.37	0.38	-2.55	1.43	0.07	0.04	-2.88	1.46	0.04	0.02
P _{trend}	0.001				0.15				0.2				0.14				0.03			
Linoleic acid																				
Q1																				
Q2	3.09	6.84	0.65	0.34	-1.03	1.31	0.43	0.83	0.76	3.57	0.83	0.9	-2.72	1.43	0.05	0.05	-2.59	1.47	0.07	0.05
Q3	5.70	8.21	0.48	0.57	0.57	1.58	0.75	0.5	-2.04	3.55	0.5	0.62	-2.08	1.43	0.14	0.11	-1.72	1.46	0.23	0.18
Q4	-3.34	10.64	0.75	0.4	0.4	2.05	0.87	0.74	1.15	3.57	0.74	0.74	-2.10	1.43	0.14	0.13	-2.34	1.47	0.11	0.12
P _{trend}	0.97				0.99				0.88				0.95				0.62			

Table 4. Multivariate-adjusted model of dietary intake of fats and semen parameters

Median for each quartile	Sperm total count				Sperm density				Total motility				DNA damage				DNA Fragmentation Index			
	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b	Beta	SE	P ^a	P ^b
Alpha-linolenic acid																				
Q1																				
Q2	3.53	6.55	0.14	0.09	0.07	1.28	0.95	0.36	0.43	3.58	0.32	0.61	-1.77	1.42	0.21	0.18	-1.83	1.46	0.21	0.16
Q3	4.27	6.65	0.55	0.31	0.33	1.27	0.31	0.11	1.17	3.58	0.04	0.06	0.36	1.41	0.8	0.81	-0.74	1.47	0.68	0.37
Q4	4.38	6.51	0.5	0.44	1.65	1.28	0.05	0.02	2.15	3.56	0.01	0.03	-2.94	1.42	0.03	0.05	-2.77	1.47	0.05	0.08
P _{trend}	0.97				0.04				0.01				0.78				0.87			
Docosapentaenoic acid (DHA)																				
Q1																				
Q2	-6.67	6.48	0.30	0.24	-0.59	1.27	0.64	0.58	3.14	3.53	0.37	0.56	0.58	1.43	0.68	0.41	0.59	1.47	0.68	0.78
Q3	-7.53	6.17	0.22	0.13	1.16	1.45	0.13	0.08	-1.49	3.36	0.65	0.52	-0.47	1.36	0.15	0.18	-0.67	1.40	0.62	0.79
Q4	-0.4	7.42	0.95	0.73	1.38	1.20	0.10	0.04	-1.10	4.04	0.78	0.72	-1.22	1.64	0.05	0.02	-1.32	1.68	0.43	0.46
P _{trend}	0.43				0.03				0.59				0.04				0.4			
Ecosapataenoic acid (EPA)																				
Q1																				
Q2	1.61	6.55	0.8	0.99	0.66	1.27	0.6	0.47	1.33	3.56	0.708	0.85	0.72	1.45	0.61	0.47	0.46	1.48	0.75	0.63
Q3	-5.85	6.55	0.37	0.33	1.97	1.27	0.12	0.15	1.13	3.56	0.246	0.31	0.03	1.45	0.98	0.83	-0.31	1.48	0.83	0.9
Q4	2.77	6.59	0.67	0.94	1.37	1.28	0.76	0.93	1.82	3.58	0.611	0.53	-0.18	1.46	0.89	0.74	-0.48	1.49	0.74	0.98
P _{trend}	0.56				0.42				0.61				0.57				0.78			

P^a: crude p- value, P^b: adjusted p-value, tests for linear trend were performed using the median value for each quartile. Multivariate model adjusted for BMI, energy intake and physical activity.

Discussion

In this cross-sectional study, it was found that higher intake of Fe, Zn, and Se were associated with higher sperm total count; Zn, ALA, DHA with semen density, Se, MUFA, and ALA with sperm total motility; DHA, Zn with DNA damage, and finally PUFA with lower percentage of DFI. Higher intake of SFA is inversely associated with sperm density. An increasing number of studies have examined the association between nutrients with semen quality (Cutillas-Tolin *et al.*, 2019, Rahimlou *et al.*, 2019, Ricci *et al.*, 2018). The findings can be compared to some extent with previous studies. Attaman J *et al.*, in a cross-sectional study, examined the association between dietary fat intake and sperm related parameters. They found that higher intake of total fat had a negative association with total sperm count and sperm concentration. Their results revealed a negative association between dietary SFA with sperm concentration and a linear association between dietary omega-3 intake with favorable sperm morphology (Attaman *et al.*, 2012). Mendiola *et al.* reported that higher intake of processed meat as one of the richest sources of SFA is associated with decreased sperm quality (Attaman *et al.*, 2012). Moreover, Safarinejad randomized 238 Iranian oligoasthenoteratospermic (OAT) men to receive a supplement containing 1840 mg EPA and DHA or placebo for 32 weeks and found that omega-3 fatty acid supplementation increased the total percentage of sperm with normal morphology (Safarinejad *et al.*, 2010), which is in line with the findings of the present study. In the present study, high intake of saturated fatty acids was associated with decreased sperm concentration. Lipids are key components of the mammal's semen. They work as structural components of semen membranes, which are precursors of numerous biologically active compounds (eicosanoids) and can be used for energy production. It has been shown that DHA is the most important spermatozoan PUFA in mammals, including man, bull, monkey, ram, and boar (Cerolini *et al.*, 2000, Poulos *et al.*, 1973). Some studies have shown that increased DHA

concentrations in semen are associated with better sperm motility (Aksoy *et al.*, 2006, Safarinejad *et al.*, 2010). The study results indicated that the participants in the higher quartile of dietary DHA intake had higher semen concentration and lower sperm DNA damage. Martínez-Soto J *et al.* in their RCT study reported that 1500 mg/day DHA supplementation for 10 weeks led to significant improvement in the seminal antioxidant status and decreased sperm DNA fragmentation (Martínez-Soto *et al.*, 2016). Safarinejad *et al.* randomized 238 Iranian OAT men to receive a supplement containing 1840 mg EPA and DHA or placebo for 32 weeks. They found that omega-3 supplementation caused a significant increase in the sperm total count, concentration, and motility (Safarinejad, 2011). Various mechanisms have been suggested for the beneficial effects of PUFA and MUFA on sperm quality. Adipose tissue depots play a significant role in the development of oxidative stress after a pro-inflammatory disorder that may alter normal reproductive pathways and sperm activity (Bachir and Jarvi, 2014, Morielli and O'Flaherty, 2015). In reality, under stress conditions, mitochondria are likely to trigger a cascade of oxidative damage in the testicular environment, since these organelles play an important role in the ROS production. MUFA and PUFA fatty acids have protective effects against increased ROS levels due to their anti-inflammatory and anti-oxidative properties (Ferramosca *et al.*, 2017, Gürler *et al.*, 2015). Evidence has suggested that these fatty acids may facilitate body fat reduction and losing weight, thereby preventing the development of obesity-related infertility; on the other, they control specific aspects of male fertility (Esmaeili *et al.*, 2015, Wathes *et al.*, 2007). PUFA supplies precursors for eicosanoid production that can attenuate the expression trends of several key enzymes (including cyclooxygenase and lipoxygenase) implicated in both prostaglandin and steroid metabolism. These fatty acids are also functional elements of cell membranes, the fluidity of which is needed to facilitate fertilization-related membrane fusion activities (Ferramosca *et al.*,

2017).

The results of the current study revealed that some minerals included Zn, Fe, and Se had beneficial effects on sperm related factors. Colagar A *et al.* in a cross-sectional study compared the concentrations of some micronutrients in semen and its association with male fertility. They found that the concentration of seminal Zn was significantly higher in the fertile subjects compared to infertile men (Colagar *et al.*, 2009). Zhao *et al.* in a meta-analysis showed that seminal plasma Zn concentrations from infertile patients were significantly lower than the control group (SMD [95% CI]: -0.64 [-1.01, -0.28]). After reviewing various studies, they reported that Zn supplementation significantly improved percentage of normal sperm morphology, sperm motility, and semen volume (Zhao *et al.*, 2016). Acute Zn depletion creates a 50% decline in Zn per ejaculate, resulting in pathozoospermia (Dissanayake *et al.*, 2010). There are several mechanisms by which Zn might interfere with sperm function. Zn as a cofactor for many metalloenzymes involved in various metabolic processes (Adewoyin *et al.*, 2017) affects phospholipases, thereby modifying the flexibility of biological membranes. It has been proposed that perhaps the elimination of Zn from the sperm cell surface disrupts the plasma membrane and plays a major role in the preparation for the completion of capacitation and the acrosome reaction. Some studies have reported that higher Zn concentration in the seminal plasma is associated with better chromatin stability (Kumar and Singh, 2016, Subhani *et al.*, 2019). It has been shown that sperm motility is most affected by Zn concentration. Zn induces stiffening the exterior dense fibers by forming disulfide bridges throughout sperm development in the epididymis, which is an important phase in the development of sperm motility; in particular, progressive motility (Kumar and Singh, 2016). Some previous findings have indicated the importance of Se and Fe in completing sperm maturation and its quality. Se is more involved in the male reproductive performance as selenoproteins. The presence of adequate Se is

important for normal spermatogenesis and sperm maturation (Ahsan *et al.*, 2014). Some previous studies have reported an inverse association between high Fe intake and sperm quality, which is not in line with the present study. Fe and copper are essential trace nutrients playing important roles in general health and fertility. However, both elements are highly toxic when accumulating in large quantities. Excessive Fe intake can stimulate the production of oxidative stress and ROS and may provide sperm membrane damage. It has been shown that either Fe excess or deficiency could lead to faulty spermatogenesis, sexual dysfunction, and oxidative damage to testicular tissue and sperm, eventually leading to impaired fertility (Ammar *et al.*, 2019, Tvrdá *et al.*, 2015).

The present study had some limitations that must be considered in interpreting the results. First, because the cross-sectional design was unable to explain any causal relationship, further prospective studies are needed to confirm the findings. Second, the plasma or seminal levels of minerals and fatty acids were not measured in the target population in this study. Finally, to record dietary intake of minerals and fatty acids a food frequency questionnaire was used that is dependent on participants' ability to remind, and their education level may increase the risk of recall bias.

Conclusion

The study results indicated that diets containing higher amounts Fe, Zn, Se, MUFA, and EPA and lower amounts of SFA were associated with more favorable semen quality parameters.

Acknowledgements

The authors would like to thank from all of the patients with contributed in this study.

Funding

There was no funding source.

Authors' contributions

Rahimlou M and Sohaei S designed the research; Rahimlou M and Nouri M contributed to acquisition of data. Sohaei S and Nouri M performed statistical analyses and interpretation of data. Sohaei S and Salehi-Sahlabadi A wrote the

manuscript. Rahimlou M and Salehi-Sahlabadi A contributed to critical revision of the manuscript. Each author has participated sufficiently in the study to take public responsibility for appropriate portions of the content.

References

- Adewoyin M, et al.** 2017. Male Infertility: The Effect of Natural Antioxidants and Phytocompounds on Seminal Oxidative Stress. *Diseases (Basel, Switzerland)*. **5 (1)**: 9.
- Afeiche M, et al.** 2013. Dairy food intake in relation to semen quality and reproductive hormone levels among physically active young men. *Human reproduction*. **28 (8)**: 2265-2275.
- Ahsan U, et al.** 2014. Role of selenium in male reproduction—A review. *Animal reproduction science*. **146 (1-2)**: 55-62.
- Aksoy Y, Aksoy H, Altinkaynak K, Aydın HR & Özkan A** 2006. Sperm fatty acid composition in subfertile men. *Prostaglandins, leukotrienes and essential fatty acids*. **75 (2)**: 75-79.
- Ammar O, Houas Z & Mehdi M** 2019. The association between iron, calcium, and oxidative stress in seminal plasma and sperm quality. *Environmental science and pollution research*. **26 (14)**: 14097-14105.
- Attaman JA, et al.** 2012. Dietary fat and semen quality among men attending a fertility clinic. *Human reproduction*. **27 (5)**: 1466-1474.
- Auger J, Kunstmann JM, Czyglik F & Jouannet P** 1995. Decline in semen quality among fertile men in Paris during the past 20 years. *New England journal of medicine*. **332 (5)**: 281-285.
- Bachir BG & Jarvi K** 2014. Infectious, inflammatory, and immunologic conditions resulting in male infertility. *Urologic clinics of North America*. **41 (1)**: 67-81.
- Cerolini S, Maldjian A, Surai P & Noble R** 2000. Viability, susceptibility to peroxidation and fatty acid composition of boar semen during liquid storage. *Animal reproduction science*. **58 (1-2)**: 99-111.
- Colagar AH, Marzony ET & Chaichi MJ** 2009. Zinc levels in seminal plasma are associated with sperm quality in fertile and infertile men. *Nutrition research*. **29 (2)**: 82-88.
- Cutillas-Tolin A, et al.** 2019. Adherence to diet quality indices in relation to semen quality and reproductive hormones in young men. *Humam reproduction*. **34 (10)**: 1866-1875.
- Dadkhah H, Kazemi A, Nasr-Isfahani M-H & Ehsanpour S** 2017. The relationship between the amount of saturated fat intake and semen quality in men. *Iranian journal of nursing and midwifery research*. **22 (1)**: 46.
- Danielewicz A, Przybyłowicz KE & Przybyłowicz M** 2018. Dietary Patterns and Poor Semen Quality Risk in Men: A Cross-Sectional Study. *Nutrients*. **10 (9)**: 1162.
- Diaz-Fontdevila M & Bustos-Obregon E** 1993. Cholesterol and polyunsaturated acid enriched diet: effect on kinetics of the acrosome reaction in rabbit spermatozoa. *Molecular reproduction and development*. **35 (2)**: 176-180.
- Dissanayake D, Wijesinghe P, Ratnasooriya W & Wimalasena S** 2010. Relationship between seminal plasma zinc and semen quality in a subfertile population. *Journal of human reproductive sciences*. **3 (3)**: 124.
- Eslamian G, Amirjannati N, Rashidkhani B, Sadeghi MR & Hekmatdoost A** 2012. Intake of food groups and idiopathic asthenozoospermia: a case-control study. *Human reproduction*. **27 (11)**: 3328-3336.
- Esmaili V, Shahverdi A, Moghadasian M & Alizadeh A** 2015. Dietary fatty acids affect semen quality: a review. *Andrology*. **3 (3)**: 450-461.
- Esmailzadeh A, et al.** 2007. Dietary patterns and markers of systemic inflammation among Iranian women. *Journal of nutrition*. **137 (4)**: 992-998.
- Fallah A, Mohammad-Hasani A & Colagar AH** 2018. Zinc is an essential element for male fertility: A review of zn roles in men's health, germination, sperm quality, and fertilization. *Journal of reproduction & infertility*. **19 (2)**: 69.
- Ferramosca A, Moscatelli N, Di Giacomo M & Zara V** 2017. Dietary fatty acids influence sperm quality and function. *Andrology*. **5 (3)**: 423-430.

- Ghaffarpour M, Houshiar-Rad A & Kianfar H** 1999. The manual for household measures, cooking yields factors and edible portion of foods. *Tehran: Nashre Olume Keshavarzy*. **7 (213)**: 42-58.
- Gürler H, Calisici O, Calisici D & Bollwein H** 2015. Effects of feeding omega-3-fatty acids on fatty acid composition and quality of bovine sperm and on antioxidative capacity of bovine seminal plasma. *Animal reproduction science*. **160**: 97-104.
- Irvine S, Cawood E, Richardson D, MacDonald E & Aitken J** 1996. Evidence of deteriorating semen quality in the United Kingdom: birth cohort study in 577 men in Scotland over 11 years. *British medical journal*. **312 (7029)**: 467-471.
- Jorgensen N, et al.** 2001. Regional differences in semen quality in Europe. *Human reproduction*. **16 (5)**: 1012-1019.
- Kumar N & Singh AK** 2016. Role of zinc in male infertility: Review of literature. *Indian journal of obstetrics and gynecology research*. **3 (2)**: 167-171.
- Künzle R, et al.** 2003. Semen quality of male smokers and nonsmokers in infertile couples. *Fertility and sterility*. **79 (2)**: 287-291.
- Lipshultz LI** 1996. "The debate continues"--the continuing debate over the possible decline in semen quality. *Fertility and sterility*. **65 (5)**: 909-911.
- Martínez-Soto JC, et al.** 2016. Dietary supplementation with docosahexaenoic acid (DHA) improves seminal antioxidant status and decreases sperm DNA fragmentation. *Systems biology in reproductive medicine*. **62 (6)**: 387-395.
- Mehra BL, et al.** 2018. Male infertility rate: a retrospective study. *Urologia journal*. **85 (1)**: 22-24.
- Morielli T & O'Flaherty C** 2015. Oxidative stress impairs function and increases redox protein modifications in human spermatozoa. *Reproduction (Cambridge, England)*. **149 (1)**: 113-123.
- Moslemi MK & Tavanbakhsh S** 2011. Selenium-vitamin E supplementation in infertile men: effects on semen parameters and pregnancy rate. *International journal of general medicine*. **4**: 99.
- Paulsen CA, Berman NG & Wang C** 1996. Data from men in greater Seattle area reveals no downward trend in semen quality: further evidence that deterioration of semen quality is not geographically uniform. *Fertility and sterility*. **65 (5)**: 1015-1020.
- Poulos A, Darin-Bennett A & White IG** 1973. The phospholipid-bound fatty acids and aldehydes of mammalian spermatozoa. *Comparative biochemistry and physiology B*. **46 (3)**: 541-549.
- Rahimlou M, Sohaei S, Nasr-Esfahani M & Nouri M** 2019. Dietary Antioxidant Intake in Relation to Semen Quality Parameters in Infertile Men: a Cross-Sectional Study. *cnr*. **8 (3)**: 229-237.
- Ricci E, et al.** 2018. Dietary habits and semen parameters: a systematic narrative review. *Andrology*. **6 (1)**: 104-116.
- Rolland M, Le Moal J, Wagner V, Royère D & De Mouzon J** 2012. Decline in semen concentration and morphology in a sample of 26 609 men close to general population between 1989 and 2005 in France. *Human reproduction*. **28 (2)**: 462-470.
- Saez Lancellotti TE, et al.** 2010. Hypercholesterolemia impaired sperm functionality in rabbits. *PloS one*. **5 (10)**: e13457.
- Safarinejad M** 2011. Effect of omega-3 polyunsaturated fatty acid supplementation on semen profile and enzymatic anti-oxidant capacity of seminal plasma in infertile men with idiopathic oligoasthenoteratospermia: a double-blind, placebo-controlled, randomised study. *Andrologia*. **43 (1)**: 38-47.
- Safarinejad MR, Hosseini SY, Dadkhah F & Asgari MA** 2010. Relationship of omega-3 and omega-6 fatty acids with semen characteristics, and anti-oxidant status of seminal plasma: a comparison between fertile and infertile men. *Clinical nutrition*. **29 (1)**: 100-105.

- Salas-Huetos A, Bullo M & Salas-Salvado J** 2017. Dietary patterns, foods and nutrients in male fertility parameters and fecundability: a systematic review of observational studies. *Human reproduction update*. **23 (4)**: 371-389.
- Sermondade N, et al.** 2013. BMI in relation to sperm count: an updated systematic review and collaborative meta-analysis. *Human reproduction update*. **19 (3)**: 221-231.
- Subhani GM, Khan NI, Akmal M, Munir MI & Javed SH** 2019. Correlation Between Semen Zinc Concentration and Semen Parameters of Infertile Men. *Annals of Punjab medical college (APMC)*. **13 (1)**: 18-22.
- Tvrda E, Peer R, Sikka SC & Agarwal A** 2015. Iron and copper in male reproduction: a double-edged sword. *Journal of assisted reproduction and genetics*. **32 (1)**: 3-16.
- Wathes DC, Abayasekara DR & Aitken RJ** 2007. Polyunsaturated fatty acids in male and female reproduction. *Biology of reproduction*. **77 (2)**: 190-201.
- Wong WY, Thomas CM, Merkus JM, Zielhuis GA & Steegers-Theunissen RP** 2000. Male factor subfertility: possible causes and the impact of nutritional factors. *Fertility and sterility*. **73 (3)**: 435-442.
- Zhao J, et al.** 2016. Zinc levels in seminal plasma and their correlation with male infertility: A systematic review and meta-analysis. *Scientific reports*. **6 (1)**: 22386.
- Zou Z, et al.** 2011. Semen quality analysis of military personnel from six geographical areas of the People's Republic of China. *Fertility and sterility*. **95 (6)**: 2018-2023. e2013.