

# Journal of

## Nutrition and Food Security

Shahid Sadoughi University of Medical Sciences School of Public Health Department of Nutrition Nutrition & Food Security Research Center



Shahid Sadoughi University of Medical Sciences School of Public Health

### eISSN: 2476-7425 pISSN: 2476-7417 JNFS 2023; 8(3): 478-485 Website: jnfs.ssu.ac.ir

## Knowledge, Attitude, and Nutritional Practice Related to COVID-19 in Students of Kurdistan University of Medical Sciences

# Ameneh Marzban; PhD<sup>1</sup>, Payam Emami; MSc<sup>2</sup>, Vahid Rahmanian; PhD<sup>3</sup>, Saeed Sherafatmanesh; MSc<sup>4</sup> & Mohsen Dowlati; PhD<sup>\*1</sup>

<sup>1</sup> Department of Health in Disasters and Emergencies, School of Health Management and Information Sciences, Iran University of Medical Sciences, Tehran, Iran; <sup>2</sup> Department of Emergency Medical Sciences, School of Paramedical Sciences, Kurdistan University of Medical Sciences, Sanandaj, Iran; <sup>3</sup> Department of Public Health, Torbat Jam Faculty of Medical Sciences, Torbat Jam, Iran; <sup>4</sup> Department of Nutrition, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

#### ARTICLE INFO

#### **ORIGINAL ARTICLE**

Article history: Received: 6 Aug 2022 Revised: 28 Sep 2022 Accepted: 19 Oct 2022

\*Corresponding author: dowlati.m@iums.ac.ir Department of Health in Disasters and Emergencies, School of Health Management and Information Sciences, Iran University of Medical Sciences, Tehran, Iran.

*Postal cod:* 1449614535 *Tel*: +98 21 88797301

#### ABSTRACT

Background: It has been known that healthy nutrition is important in increasing immunity and reducing diseases. Lack of some nutrients can lead to a decrease in the function of the immune system and, as a result, an increase in the probability of infections or their exacerbation. This study was conducted with the aim of investigating knowledge, attitude and nutritional practice related to COVID-19 in the students of Kurdistan University of Medical Sciences. Methods: The present descriptive and analytical study was conducted in 2022 on 350 students of Kurdistan University of Medical Sciences. The research tools included a demographic checklist and a questionnaire of knowledge, attitude and nutritional practice related to COVID-19, used in a similar study. The collected data were analyzed by SPSS 24 software using descriptive statistics and analysis of variance, independent t-test, and Pearson's correlation coefficient. Results: The mean age of the subjects was 22.15±4.98 year. The mean scores of knowledge, attitude, and nutritional behavior of students of Kurdistan University of Medical Sciences regarding COVID-19 were 23.17±4.17, 18.04±3.29, and  $41.64 \pm 7.16$ , respectively. The results showed that there was a statistically significant relationship between the knowledge score with the educational level and history of infection with COVID-19 and the nutritional practice score with the variable of history of infection with COVID-19 (P < 0.05). Also, there was a positive and significant correlation between Knowledge, attitude, and practice. Conclusion: According to the results of the study, the studied students had a relatively good knowledge, attitude, and practice towards nutritional issues

This paper should be cited as: Marzban A, Emami P, Rahmanian V, Sherafatmanesh S, Dowlati M. Knowledge, Attitude, and Nutritional Practice Related to COVID-19 in Students of Kurdistan University of Medical Sciences. Journal of Nutrition and Food Security (JNFS), 2023; 8(3): 478-485.

#### related to COVID-19; however, there is still room for improvement.

Keywords: Knowledge; Attitude; Behavior; Nutrition; Students; COVID-19

#### Introduction

Noronaviruses are a large family of viruses that, in addition to mild to severe infections of the respiratory tract, can also involve other vital organs of the body and in acute cases cause digestive problems such as diarrhea, coagulation disorders, heart problems and kidney failure (Mitchell, 2020). There are hundreds of different types of corona virus, most of which are found in animals such as pigs, camels, bats, and cats, and seven of them are known to cause disease in humans (Seah and Agrawal, 2020). In December 2019, an outbreak of SARS-CoV-2 was reported from the city of Wuhan, China, and caused the disease of COVID-19 (Marzban et al., 2022). The World Health Organization (WHO) announced it a global pandemic in 2020 (Salehi et al., 2022). Approximately two-thirds of the identified patient cases were linked to a seafood market in Wuhan, where live aquatic animals such as civets and raccoons were sold (Chen et al., 2020). COVID-19 is a highly contagious infectious disease that is transmitted from person to person directly through respiratory droplets of infected people when they sneeze or cough (Morawska and Cao, 2020). The behavior of people in the society has a direct effect on the control and prevention of COVID-19 disease, and the transmission process of this disease can be affected by public health behaviors (La Torre et al., 2021). In the absence of a specific treatment for this new virus, finding alternative methods can play an important role in preventing, controlling, multiplying, and spreading the virus. Effective nonpharmacological interventions combined with promoting a healthy lifestyle and beneficial dietary patterns can improve overall health and reduce the risk of infection due to SARS-CoV-2 (Hibino and Hayashida, 2022). Following a balanced and healthy diet that contains all the essential nutrients is very important to maintain health (Misra, 2020). Macronutrients (fats, proteins, and carbohydrates) and micronutrients (vitamins and minerals) promote healthy immune responses and for many years, the importance of healthy nutrition has been known in increasing immunity and reducing diseases (Taghdir et al., 2020). Many studies have shown that a lack of certain nutrients such as zinc, selenium, iron, vitamins E, D, C, B, A, proteins, and omega-3 fatty acids can lead to a decrease in the immune system and an increase in the probability of infections and or their aggravation (Alsan et al., 2020, Luo et al., 2021). On the other hand, antiinflammatory nutrients such as beta-carotene, vitamin C, vitamin E, vitamin B3, probiotics, and polyphenolic compounds can strengthen the function of the immune system through the production and activity of antimicrobial proteins and increasing the activity of neutrophils and macrophages (Calder et al., 2020, Martineau et al., 2017). Therefore, the anti-inflammatory strategy by nutrients can be considered as a suitable option in the management of the COVID-19 disease (Conti et al., 2020, Kritas et al., 2020). Based on studies, increasing people's knowledge about health issues has a very favorable effect on improving health status and leads to healthier nutritional practices (Jahed et al., 2012, Marzban et al., 2022). The role of medical students in raising knowledge during public health emergencies has long been proven, and medical students with higher levels of health literacy may act as public role models for adopting preventive health behaviors against COVID-19 (Martin et al., 2022). It is required to design

strategies to encourage more communities to follow health standards. In addition, lifestyle and dietary modification can improve overall health status and prevent diseases other than those associated with COVID-19 (Hibino and Hayashida, 2022). This information may also be used by medical universities to improve curricula to prepare for future disease outbreaks (Adli *et al.*, 2022). Therefore, considering the importance of students' knowledge and attitude towards food safety during the COVID-19 and lack of related studies among medical students, the present study aimed to investigate the knowledge, attitude, and nutritional practice related to COVID-19 in students of Kurdistan University of Medical Sciences.

#### **Materials and Methods**

Study design and participants: This descriptiveanalytical study was carried out in 2022. The study population consisted of all students of Kurdistan University of Medical Sciences. The sample size was determined at 350 people using similar studies (Mansoorian *et al.*, 2021) through the following formula, taking into account the confidence level of 95% and Z=1.96.

$$n = \frac{z^2 p(1-p)}{d^2}$$

*Measurements*: The demographic information asked from the participants included age, gender, educational level, economic status, college, marital status, and number of family members. The questionnaire included three questions about the history of chronic diseases such as diabetes, blood pressure, etc., as well as the history of COVID-19.

The Questionnaires were completed online in the press line space, and prior to completion, necessary arrangements were made to obtain informed consent through virtual space and phone calls. Questionnaire link was sent to each person via email or social networks such as WhatsApp or Telegram.

The research tool of knowledge, attitude, and nutritional practice questionnaire related to the COVID-19 was examined in the study of Marzban and Mansoorian (Mansoorian *et al.*, 2021, Marzban *et al.*, 2022) in Yazd City, and its validity and reliability were confirmed. Cronbach's alpha in three parts of knowledge, belief, and nutritional behavior about COVID-19, were obtained 0.82, 0.79, and 0.83, respectively.

The questionnaire included 20 questions related to people's nutritional knowledge about COVID-19, which were based on a 3-point Likert scale, including 'correct', 'incorrect', and 'I do not know'. The 'correct' answer was scored 2, and the answer 'I don't know' was scored 1 and the 'incorrect' was scored zero. The obtained score of each person in the knowledge section was in the range of 0-40.

The questionnaire included 13 questions related to the nutritional attitude of people regarding the COVID-19, and the answers were based on a threepoint Likert scale, including 'I agree', 'I have no opinion', and 'I disagree', graded from 1 to 3. The obtained score of each person in the nutritional attitude section was in the range of 13-39.

The questionnaire included 20 questions related to the nutritional practice of people regarding the COVID-19, and the answers were based on a threepoint Likert scale, including 'always', 'sometimes' and 'never'. The answers were scored from 1 to 3. The obtained score of each person in the nutritional practice section was in the range of 20-60.

The inclusion criteria were being a student of Kurdistan University of Medical Sciences and participating in the program until the end of the study. The exclusion criteria were unwillingness to cooperate or an incomplete questionnaire. Before starting the research, the conditions were explained to them and finally those who wanted to participate in the study were included in the study.

*Data analysis:* The collected data were entered into SPSS 24 software and analyzed with Pearson's correlation coefficient tests, descriptive statistics, chi-square, t-test, and analysis of variance.

*Ethical considerations:* It should be mentioned that the present study is taken from a research project at Kurdistan University of Medical Sciences, with a code of ethics approved by the Ethics Committee IR.MUK.REC.1401.116.

#### **Results**

The mean age of the participants was  $22.15\pm4.98$  year. The results showed that among the 350 students, the majority (54.85%) were female, and

36.28% of them were in the age range of 21-24 year. Also, 55.16% of them were undergraduate students, 23.71% were from Nursing-Midwifery Faculty, 56.57% of the students had a moderate economic status, and 66% of the participants were married. Moreover, 53.14% of the students had been infected with COVID-19 and 90.58% had no history of chronic diseases. General information about the demographic characteristics of the students is given in **Table 1**.

The mean scores of knowledge, attitude, and nutritional practice of Kurdistan University of Medical Sciences students regarding COVID-19 were  $23.17\pm4.17$ ,  $18.04\pm3.29$ , and  $41.64\pm7.16$ ,

respectively.

According to **Table 2**, the results showed that there was a statistically significant relationship between the knowledge score with education level and the history of being infected with COVID-19.

**Table 3** reveals that the most important source ofinformation for people regarding COVID-19 wascyberspace, and people obtained the leastinformation through television and radio.

Results show a positive and significant correlation between the variables of knowledge, attitude, and practice, and the correlation coefficient between attitude and practice was greater.

Table 1. Frequency distribution of demographic variables in Kurdistan University of Medical Sciences students.

| Variables                               | Description                    | N   | %     |
|---|--------------------------------|-----|-------|
| Gender                                  | Male                           | 158 | 45.15 |
|   | Female                         | 192 | 54.85 |
| Age (year)                              | <21                            | 112 | 32.00 |
|   | 21-2,4                         | 127 | 36.28 |
|   | >24                            | 111 | 31.72 |
| Level of education                      | Bachelor                       | 193 | 55.16 |
|   | Master's degree                | 89  | 25.42 |
|   | PhD and professional doctorate | 68  | 19.42 |
| Faculty                                 | Medical                        | 65  | 18.57 |
|   | Paramedical                    | 62  | 17.71 |
|   | Dental                         | 63  | 18.00 |
|   | Health                         | 78  | 22.28 |
|   | Nursing and Midwifery          | 83  | 23.71 |
| Economic status of the family           | Good                           | 84  | 24.00 |
| Economic status of the family           | Moderate                       | 198 | 56.57 |
|   | Weak                           | 68  | 19.42 |
| Marital status                          | Single                         | 231 | 66.00 |
|   | Married                        | 119 | 34.00 |
| A person's history of COVID -19         | Yes                            | 186 | 53.14 |
| A person's history of COVID -19         | No                             | 164 | 46.86 |
| A person's history of chronic diseases  | Yes                            | 317 | 90.58 |
| A person's instory of enfolite diseases | No                             | 33  | 9.42  |

 Table 2. Mean score of knowledge, attitude, and practice of Kurdistan University of Medical Sciences students according to demographic variables.

| Variables            | Degenintian | knowl | knowledge |       | Attitude |       | Practice |  |
|----------------------|-------------|-------|-----------|-------|----------|-------|----------|--|
|                      | Description | Mean  | SD        | Mean  | SD       | Mean  | SD       |  |
| Gender               | Male        | 23.84 | 4.25      | 18.69 | 3.57     | 41.01 | 7.95     |  |
|                      | Female      | 23.64 | 4.19      | 18.07 | 3.09     | 41.33 | 7.35     |  |
| P-value <sup>a</sup> |             | 0.2   | .4        | 0.2   | 21       | 0.1   | 7        |  |
| Age (year)           | <21         | 23.57 | 4.12      | 18.69 | 3.55     | 41.12 | 7.14     |  |
|                      | 21-24       | 23.15 | 4.19      | 18.07 | 3.67     | 41.39 | 7.06     |  |
|                      | >24         | 23.17 | 4.78      | 18.67 | 3.33     | 41.65 | 7.31     |  |

| Variables                     | Description           | knowl     | knowledge |       | Attitude |       | Practice |  |
|-------------------------------|-----------------------|-----------|-----------|-------|----------|-------|----------|--|
| v al lables                   |                       | Mean      | SD        | Mean  | SD       | Mean  | SD       |  |
| P-value <sup>b</sup>          |                       | 0.3       | 32        | 0.30  |          | 0.27  |          |  |
| Level of education            | Bachelor              | 19.24     | 4.31      | 18.24 | 3.60     | 41.03 | 7.28     |  |
|                               | Master's degree       | 26.25     | 4.57      | 18.54 | 3.95     | 41.97 | 7.09     |  |
|                               | PhD and professional  | 34.12     | 4.87      | 18.03 | 3.54     | 41.52 | 7.16     |  |
|                               | doctorate             |           |           |       |          |       |          |  |
| P-value <sup>b</sup>          |                       | <0.0      |           | 0.25  |          | 0.18  |          |  |
| Faculty                       | Medical               | 23.75     | 4.54      | 18.17 | 3.01     | 41.34 | 7.49     |  |
| Faculty                       | Paramedical           | 23.64     | 4.13      | 18.32 | 3.44     | 41.05 | 7.06     |  |
|                               | Dental                | 23.84     | 4.03      | 18.94 | 3.09     | 41.26 | 7.66     |  |
|                               | Health                | 23.34     | 4.06      | 18.24 | 3.88     | 41.08 | 7.37     |  |
|                               | Nursing and Midwifery | 23.15     | 4.11      | 18.06 | 3.91     | 41.21 | 7.13     |  |
| P-value <sup>b</sup>          |                       | 0.3       | 85        | 0.34  |          | 0.21  |          |  |
| Economic status of the family | Good                  | 23.17     | 4.07      | 18.22 | 3.14     | 40.29 | 7.01     |  |
|                               | Moderate              | 23.16     | 4.69      | 18.57 | 3.15     | 41.35 | 7.33     |  |
|                               | Weak                  | 22.98     | 4.08      | 18.35 | 3.06     | 41.87 | 7.79     |  |
| P-value <sup>b</sup>          |                       | 0.14 0.19 |           | .9    | 0.16     |       |          |  |
| Marital status                | Single                | 23.66     | 4.88      | 18.54 | 3.19     | 40.18 | 7.46     |  |
|                               | Married               | 23.19     | 4.60      | 18.09 | 3.45     | 40.09 | 7.69     |  |
| P-value <sup>a</sup>          |                       | 0.20 0.17 |           | 7     | 0.24     |       |          |  |
| A person's history of COVID - | Yes                   | 37.12     | 4.11      | 18.67 | 3.67     | 41.11 | 7.37     |  |
| 19                            | No                    | 23.11     | 4.05      | 18.02 | 3.12     | 41.32 | 7.19     |  |
| P-value <sup>a</sup>          |                       | < 0.001   |           | 0.21  |          | 0.15  |          |  |
| A person's history of chronic | Yes                   | 23.11     | 4.66      | 18.33 | 3.68     | 41.06 | 7.42     |  |
| diseases                      | No                    | 23.17     | 4.51      | 18.04 | 3.49     | 41.11 | 7.56     |  |
| P-value <sup>a</sup>          |                       | 0.23      |           | 0.31  |          | 0.25  |          |  |

 Table 2. Mean score of knowledge, attitude, and practice of Kurdistan University of Medical Sciences students according to demographic variables.

<sup>a</sup>: Independent Samples t-test, <sup>b</sup>: ANOVA test.

# Table 3. The frequency of the source of information about the Covid-19 in the students of Kurdistan University of Medical Sciences.

| Sources                           | Ν   | %     |
|-----------------------------------|-----|-------|
| Cyberspace and internet           | 75  | 21.42 |
| TV and radio                      | 41  | 11.71 |
| Doctor and health center staff    | 56  | 16.00 |
| Friends and family                | 58  | 16.57 |
| Book and magazine                 | 52  | 14.85 |
| University professors and courses | 68  | 19.42 |
| Total                             | 350 | 100   |

#### Discussion

One of the most important measures in maintaining health and preventing infectious diseases is to have proper and basic nutrition and enough sleep. This study was conducted with the aim of investigating the knowledge, attitude, and nutritional practice related to COVID-19 among the students of Kurdistan University of Medical

#### Sciences.

The mean age of the participants in the study was  $22.15\pm4.98$  year. In the study of Marzban (Marzban *et al.*, 2022) and Mansourian (Mansoorian *et al.*, 2021), the mean age of the subjects was  $36.57\pm5.24$  and  $37.30\pm11.30$  year. In Luo's study (Luo *et al.*, 2021) among Chinese people, the mean age of people was  $24.09\pm9.14$  year and 54.85% of

participants were female. In Alhaj's study (Alhaj et al., 2021) in Jordan, 69.5% of the participants were female. However, in the study of Marzban (Marzban et al., 2022) and Mansourian (Mansoorian et al., 2021), 56.90% and 52.70% of the participants were male. Women seem to be more likely to respond to diet and health surveys, since they are usually responsible for the diet and nutrition of families (Al-Sheyab et al., 2019). In the present study, 53.14% of the students reported a history of being infected with COVID-19; however, in Marzban's study, 61.62% of the them were not infected with COVID-19. The difference in the number of infected cases can be attributed to the spread of the omicron strain, which involved most of people in the community due to its greater contagion.

Marzban A, et al.

The mean score of nutritional knowledge of Kurdistan University of Medical Sciences students was 23.17±4.17 in relation to COVID-19. The mean nutritional knowledge score of the people of Yazd city was reported 21.35±4.54 in Marzban's study (Marzban et al., 2022). In Mansourian's study (Mansoorian et al., 2021), which investigated nutritional knowledge, attitude, and practice of the people of Gonabad city, the mean nutritional knowledge score of people was 9.70±1.40. In Luo's study (Luo et al., 2021), examining knowledge, attitude, and practice of Chinese people in the field of food safety and nutrition during the COVID-19, the mean knowledge score of people was reported 5.20±1.60. Knowledge of nutrition and diet is an integral part of health knowledge. In China, the "Healthy China 2030" plan with the aim of promoting and improving food safety and the nutritional status of Chinese residents is considered as an important content and goal (Luo et al., 2021). Knowledge is a key tool for patients and their families to have a correct understanding of the COVID-19, and not having enough knowledge can disrupt the process of modifying food habits and behaviors. Nutrition knowledge and related skills can also enable people to make appropriate food choices and improve their metabolic status and quality of life. There has been no extensive research on the relationship between nutritional knowledge JNFS | Vol (8) | Issue (3) | August 2023

and food intake in patients with COVID-19 due to the new nature of this disease; however, it has been proven that nutrition has an important role in viral and infectious diseases and proper nutrition can improve the immune system (Al-Badayneh, 2021, Sami *et al.*, 2021).

The mean score of nutritional attitude of the students towards COVID-19 was 18.04±3.29. The mean score of attitude in the study of Marzban 2022) (Marzban et al., and Mansourian (Mansoorian et al., 2021) was 20.74±3.58 and 14.3±3.5. With the development of society and improvement of living standards, people's attention to food safety and nutrition has increased. Food behavior is the result of the interaction between human knowledge and attitude. Knowledge refers to the knowledge of things and processes that specialists non-specialists. distinguish from Attitudes are people's opinions that are influenced by knowledge and lead to certain actions (practice). Therefore, behaviors vary from simple behaviors such as chewing food to complex ones such as food preparation, social etiquette, and policy making. Lack of health literacy is associated with poor health outcomes. However, people may have necessary knowledge about nutrition, but the correct attitude has not been formed in them, and this knowledge will not play a role in improving people's health.

The mean score of people's nutritional practice in relation to COVID-19 is 41.64±7.16. The mean score of nutritional practice in the study of Marzban (Marzban et al., 2022) and Mansourian (Mansoorian et al., 2021) was 38.99±7.39 and 29.6±4.2. The mean practice score of Chinese people in Luo's study (Luo et al., 2021) was 3.7±1.0. In this study, many Chinese residents changed their food safety behaviors and dietary patterns, including increasing their consumption of vegetables, fruits, and water and decreasing their consumption of sugary drinks and snacks. However, in Alhaj's study (Alhaj et al., 2021) in Jordan, the participants did not pay much attention to their diet during the COVID-19. Regarding nutrition, two important questions may be raised: "What food should be consumed to strengthen the immune system?" "How can we consume food to prevent

contamination?" In response to the first question, it is recommended to avoid consuming open or bulk foods, traditional dairy products, cakes, cookies, dumplings, and nuts in centers that are served in open or bulk form. Protein materials such as eggs, all kinds of meats, kebabs, and steaks should be well cooked. Eating raw food should be avoided and cooked food should not come in contact with raw food. Corona virus is destroyed by heat, industrial packaging can help prevent this virus, and when buying bread, it should be put directly to the buyer's clean hand using clean gloves. Intermediaries such as metal surfaces of bakeries can be dangerous; therefore, surfaces should be frequently disinfected and the contamination of intermediaries in this way should be reduced as much as possible. The best and most important way to strengthen the immune system is to use food rich in vitamins. White blood cells need vitamins to provide enough energy to fight viruses, so it is important to identify the role of different vitamins in the body's immune system and food ingredients containing these vitamins (Ramezani and Amirpour, 2020).

The results showed that there is a statistically significant relationship between the knowledge score with the level of education and the history of COVID-19 infection. This finding was consistent with the results of Marzban (Marzban et al., 2022) and Luo (Luo et al., 2021) studies . However, in Mansourian's study (Mansoorian et al., 2021), no significant relationship was observed between demographic variables and knowledge scores. Higher education leads to more information about viral diseases, and since the study population was medical students, with the advancement of education level, they take more courses on infectious diseases and nutrition. Also, the history of the disease causes a person to be more sensitive about ways of recovery and prevention.

#### Conclusion

484

According to the results of the study, the studied students had a relatively good knowledge, attitude, and practice towards the nutritional issues related to the COVID-19; however, there is still room for improvement. Research has shown that nutrition knowledge interventions can effectively improve people's understanding of relevant knowledge and, as a result, create positive changes in attitudes and practices. Public education on healthy nutrition should be promoted as a normal subject and widely implemented in various units and departments at the public level. Virtual space, which occupies an important part of people's lives, should be considered in this case.

#### Acknowledgment

The authors would like to thank Kurdistan University of Medical Sciences for its financial support, as well as students of Kurdistan for their participation in the study.

#### Authors' contributions

Marzban A and Dowlati M were involved in designing and supervising the survey. Emami P and Sherafatmanesh S were involved in designing the study, data collecting, and data analyzing. Rahmanian V and Marzban A participated in data analysis. Marzban A, Emami P and Dowlati M participated in writing the manuscript. All authors critically reviewed the manuscript and approved the final version submitted for publication.

#### **Conflict of interest**

The authors declared no conflict of interest.

#### References

- Adli I, et al. 2022. Knowledge, attitude, and practice related to the COVID-19 pandemic among undergraduate medical students in Indonesia: A nationwide cross-sectional study. *PloS one.* **17** (1): e0262827.
- Al-Badayneh G 2021. The Impact of Strategic Awareness on Enhancing Organizational Immunity System: An Applied Study on Jordanian Food Manufacturing Companies. International review of management and marketing. 11 (2): 47-58.
- Al-Sheyab NA, Alomari MA, Hayajneh AA &
  Shah S 2019. Attitudes and perceived barriers toward healthy lifestyle behaviors in Jordanian adolescents: a developing country perspective. *Adolescent health, medicine and therapeutics.* 10: 39.

DOI: 10.18502/jnfs.v8i3.13295

- Alhaj OA, Al-Sayyed HF, AlRasheed MM & Jahrami H 2021. Appraisal survey of the knowledge, attitudes, and behaviors of Jordanian society toward diet and nutrition during COVID-19 era. *Journal of public health research.* **10** (4): jphr. 2021.2381.
- Alsan M, Stantcheva S, Yang D & Cutler D 2020. Disparities in coronavirus 2019 reported incidence, knowledge, and behavior among US adults. *JAMA network open.* 3 (6): e2012403e2012403.
- Calder PC, Carr AC, Gombart AF & Eggersdorfer M 2020. Optimal nutritional status for a well-functioning immune system is an important factor to protect against viral infections. *Nutrients.* **12** (4): 1181.
- Chen Y, Liu Q & Guo D 2020. Emerging coronaviruses: genome structure, replication, and pathogenesis. *Journal of medical virology*. 92 (4): 418-423.
- Conti P, et al. 2020. Induction of pro-inflammatory cytokines (IL-1 and IL-6) and lung inflammation by Coronavirus-19 (COVI-19 or SARS-CoV-2): anti-inflammatory strategies. *Journal of biological regulators and homeostatic agents*. 34 (2): 327-331.
- Hibino S & Hayashida K 2022. Modifiable Host Factors for the Prevention and Treatment of COVID-19: Diet and Lifestyle/Diet and Lifestyle Factors in the Prevention of COVID-19. *Nutrients.* **14** (9): 1876.
- Jahed G, GolestaniFar H, Ghodsi R & Mohammadi M 2012. The knowledge and attitude of students in relation with health and food safety at Tehran University of Medical Sciences. *Journal of research and health.* 2 (1): 72-80.
- Kritas S, et al. 2020. Mast cells contribute to coronavirus-induced inflammation: new antiinflammatory strategy. *Journal of biological regulators and homeostatic agents.* 34 (1): 9-14.
- La Torre G, et al. 2021. How much do young Italians know about COVID-19 and what are their attitudes toward SARS-CoV-2? results of a crosssectional study. *Disaster medicine and public health preparedness.* **15** (1): e15-e21.

- Luo Y, et al. 2021. Investigation on knowledge, attitudes and practices about food safety and nutrition in the China during the epidemic of corona virus disease 2019. *Public health nutrition.* 24 (2): 267-274.
- Mansoorian M, Noori R, Khosravan S, Tabatabaeizadeh S-A & Khajavian N 2021. Nutritional knowledge, attitude and behaviours regarding Coronavirus Disease 2019 among residents of Gonabad, Iran. *Public health nutrition.* 24 (5): 1088-1094.
- Martin A, et al. 2022. A rapid systematic review exploring the involvement of medical students in pandemics and other global health emergencies. *Disaster medicine and public health preparedness.* 16 (1): 360-372.
- Martineau AR, et al. 2017. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *British medical journal.* 356.
- Marzban A, et al. 2022. Nutritional Knowledge, Attitude, and Practices Related to COVID-19 in People of Yazd, 2021. *Journal of nutrition and food security*. **7** (1): 22-29.
- Misra A 2020. Balanced nutrition is needed in times of COVID19 epidemic in India: a Call for Action for all nutritionists and physicians. *Diabetes & metabolic syndrome*. **14** (6): 1747-1750.
- Mitchell EP 2020. Corona virus: Global pandemic causing world-wide shutdown. *Journal of the national medical association*. **112** (2): 113.
- Morawska L & Cao J 2020. Airborne transmission of SARS-CoV-2: The world should face the reality. *Environment international*. **139**: 105730.
- Ramezani A & Amirpour M 2020. Nutritional Care in the Prevention and Treatment of Coronavirus Disease 2019: A Simple Overview. *Journal of health research in community.* 6 (1): 74-82.
- Salehi A, Salmani F, Norozi E, Sadighara P & Zeinali T 2022. Knowledge, attitudes and practices of Iranian people about food safety and hygiene during covid-19 pandemic. *BMC public health.* 22 (1): 1-9.

- Sami R, Alshehry G, Elgarni E & Helal M 2021. Saudi community care awareness food facts, nutrients, immune system and COVID-19 prevention in Taif city among different age categories. *African journal of food, agriculture, nutrition and development.* **21** (1): 17213-17233.
- Seah I & Agrawal R 2020. Can the coronavirus disease 2019 (COVID-19) affect the eyes? A

review of coronaviruses and ocular implications in humans and animals. *Ocular immunology and inflammation.* **28** (**3**): 391-395.

Taghdir M, Sepandi M, Abbaszadeh S & Parastouei K 2020. A review on some Nutrition-Based Interventions in Covid-19. *Journal of military medicine*. 22 (2): 169-176.