



## Application of Dispersive Liquid-Liquid Microextraction to Determine Aflatoxin B1 in Tomato Paste Samples

Vahid Safavizadeh; MSc<sup>1</sup>, Pouya Arabkhani; PhD<sup>2</sup>, Mozhgan Mojkar; MSc<sup>3</sup>,  
Darya Shyrina; PhD<sup>4</sup> & Mahboob Nemati; PhD\*<sup>5</sup>

<sup>1</sup> Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran.

<sup>2</sup> Department of Pharmaceutical and Food Control, School of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

<sup>3</sup> Department of Human Ecology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

<sup>4</sup> Department of Technology Processing of Fruits, Vegetables and Milk, Kharkov State University of Food Technology and Trade, Kharkiv, Ukraine.

<sup>5</sup> Halal Research Center of IRI, FDA, Tehran, Iran.

### ARTICLE INFO

#### ORIGINAL ARTICLE

##### Article history:

Received: 9 Jun 2019

Revised: 21 Jun 2020

Accepted: 24 Dec 2019

##### \*Corresponding author:

nematim@tbzmed.ac.ir

Halal Research Center of  
IRI, FDA, Tehran, Iran.

Postal code: 5166414766

Tel: +98 09142353619

### ABSTRACT

**Background:** Aflatoxin B1 is a secondary toxic metabolite produced by several aspergillus species. This study was conducted to determine the presence of aflatoxin in tomato paste samples. **Methods:** A total of 30 tomato pastes were analyzed for Aflatoxin B1 contamination via HPLC-FLD. Analyte extraction was done by dispersive liquid-liquid microextraction. **Results:** The limit of detection (LOD) and limit of quantitation (LOQ) were 0.14 and 0.44 µg/kg, respectively. The recovery rates ranged from 91 to 94%. The findings showed all samples were contaminated with Aflatoxin B1 and the average concentration was  $1.1 \pm 0.02$  µg/kg. The amount of aflatoxin B1 in 6 samples was higher than the limit set by the European Union. **Conclusion:** The proposed method was successfully applied to the analysis of tomato paste samples with quantitative results. The main advantages of the developed method include its simplicity in operation, rapid achievement of a very high sample, and low cost.

**Keywords:** Mycotoxins; HPLC; Aflatoxin B1; Iran

### Introduction

Mycotoxins are the secondary metabolites produced by different types of fungi (Flores-Flores *et al.*, 2015). Their presence in food depends on a variety of factors, such as temperature, humidity, storage, and Processing (Rawat, 2015). Aflatoxin is one of the most toxic known mycotoxins produced by genus *Aspergillus*, mostly *aspergillus flavus* and *aspergillus parasiticus* species (Yu *et al.*, 2004).

Aflatoxin B1 (AFB1), aflatoxin B2 (AFB2), aflatoxin G1 (AFG1), and aflatoxin G2 (AFG2) are

the most important types of aflatoxin. Furthermore, aflatoxin M1 (AFM1) and aflatoxin M2 (AFM2) are metabolites of AFB1 and AFB2 in the livestock body (Kumar *et al.*, 2017), which are generally found in a wide range of foods such as grains, pistachio, milk, and nuts (Alshannaq and Yu, 2017). Aflatoxins are highly toxic, mutagenic, teratogenic, and carcinogenic compounds that are known as the causative agents of liver carcinogenesis. The most toxic aflatoxin is AFB1 that was determined by the

International Agency for Research on Cancer (IARC) as group 1 carcinogen types (Abrar *et al.*, 2013).

The allowed limits of AFB1 and total aflatoxin are 2 and 4 µg/kg in food products in the European Union, respectively (Authority, 2007, Kilicel *et al.*, 2017). Various methods were introduced to detect aflatoxins, such as ELISA and HPLC using mass detection and several other methods. In this regard, HPLC with fluorescence detection (FLD) is one of the most commonly used methods to detect aflatoxin (Espinosa-Calderón *et al.*, 2011).

The dispersive liquid-liquid microextraction (DLLME) technique was introduced in 2006, which is a quick, low-cost, and low-solvent consumption method. This method is used to extract a small amount of the material from aqueous samples. This technique is based on injecting a small amount of a solvent extraction agent (like C2Cl4) and a dispersed solvent (like acetone) into an aqueous solution. The resulting solution is then centrifuged to separate the organic and aqueous layers. This process is useful in extracting organic compounds from aqueous samples (Berijani *et al.*, 2006, Rezaee *et al.*, 2006).

In previous studies over analysis of mycotoxins using the development and rapid sample preparation methods, the suitability of DLLME was investigated for sensitive determination of aflatoxins in samples. HPLC\_FLD was used as a determination technique (Campone *et al.*, 2011).

In this study, the aim was to measure the amount of aflatoxin extracted using DLLME method in tomato paste samples collected from the northwest of Iran. Later, the amount of aflatoxin contamination in Iran was compared with the international standards.

## Materials and Methods

**Chemicals and reagents:** Sodium chloride (NaCl), n-hexane, chloroform (CHCl<sub>3</sub>), and Formic acid (CH<sub>2</sub>O<sub>2</sub>) included high purity LC grade provided by Merck (Darmstadt, Germany). The stock solution of AFB1 at a concentration of 1000.0 ng/ml was supplied by Sigma (St.Louis, MO, USA). Methanol and acetonitrile were prepared from Duksan Pure Chemicals (Kyungkido, Korea), deionized extra pure water was also purchased from Shahid Ghazi pharmaceutical company (Tabriz-Iran). To construct

a calibration curve, 6 dilutions of AFB1 standards were prepared from 0.6, 1.25, 2.5, 5, 10, and 20 ng/ml by adding appropriate volumes of methanol to obtain these concentrations. The Stock solution of AFB1 was kept at 4 °C in the dark.

**Equipment and chromatographic conditions:** The HPLC analyses were carried out in a KNAUER HPLC system with a K-1000 pump (Berlin, Germany) coupled with a fluorescence detector. Biotech 2003 degasser (United State) and the analytical column included C18 column 4.6x250 mm (Knauer, Berlin, Germany).

The mobile phase was acetonitrile: methanol: water (24:25:51), de-aerated by an ultrasonic bath for 10 min at a flow rate of 1.5 ml/min in isocratic mode. The fluorescence detector was set at wavelengths of 373 and 450 nm for excitation and emission, respectively.

**Method of validation:** The developed analytical method was validated based on repeatability (intra-day precision) and reproducibility (inter-day precision), which is also called RSD, linearity, accuracy (as recoveries), selectivity, as well as limits of detection (LODs) and quantification (LOQs). The LOD was calculated as the concentration of analyte, whose peak area was three times the area of the noise of a blank sample ( $S/N \geq 3$ ). LOQ was calculated by taking three replicates of the detection limits when  $S/N \geq 10$ . The acceptable value for RSD is below 20% and the acceptable range for recovery is 70-120% (Fontana *et al.*, 2016). In measuring linearity, a value of higher than 0.99 is acceptable.

**Tomato paste sampling:** Tomato paste samples were collected from different stores in 2019 within two months. A total of 30 samples were collected from different brands, including 10 famous Iranian brands, 12 non-famous Iranian brands, 5 non-Iranian brands, and 3 homemade tomato pastes.

**Extraction procedure:** The extraction procedure of analyte from tomato paste samples was carried out according to a previous study method (Amirkhizi *et al.*, 2015) after applying some modifications. To this end, 5 g well-homogenized tomato paste samples were mixed with 0.1 g NaCl and 20 ml acetonitrile-

water (80:20, V/V). Later, they were blended by a magnetic stirrer for 10 minutes; 1 g diatomaceous earth was added and mixed for 5 min and the mixture was clarified with Whatman filter paper No.1 (Whatman Inc., Clifton, NJ, USA).

In the next stage, 3 ml of n-Hexane was added to 10 ml of the extracted solution and shortly shaken by Vortex. The n-Hexane was separated by centrifugation for 1 min and cleaned up by DLLME (Amirkhizi *et al.*, 2015).

*Dispersive liquid-liquid microextraction procedure:* A volume of 260 µl chloroform (extracting solvent) was added to 1 ml of the obtained extract (dispenser solvent) and the pH was adjusted to about 5.8 by formic acid. The mixture was quickly injected into a 10 ml centrifuge test tube with a conical bottom containing 2.5 ml of distilled water. It was then vortexed for a few seconds to obtain a stable and cloudy solution; so, AFB1 was entrapped into the fine CHCl<sub>3</sub> droplets. This ternary component solution was centrifuged at 5000 RPM for 3 minutes after the aqueous upper layer was removed. The precipitated chloroform was transferred to a small tube by a 220 µl sampler and dried at a mild flow of nitrogen gas. The residue was re-dissolved in 20 µl methanol and then injected into the HPLC system (Afzali *et al.*, 2012, Amirkhizi *et al.*, 2015).

### Results

*Method validation:* In order to verify the linearity, six concentrations of the standard B1 0.6, 1.25, 2.5, 5, 10, and 20 ng/ml were injected into the device in three replications and obtained with a calibration curve  $R^2 = 0.9974$ , which is acceptable. The LOD and LOQ were 0.14 and 0.44 µg/kg, respectively. The recovery and RSD rates were acceptable (**Table 1**).

*Testing the method for evaluating the natural occurrence of AFB1 in tomato pastes:*

The analytical results of AFB1 in samples are reported in **Table 2**. AFB1 was detected in all samples and quantified in 14 samples. In this study, concentration ranged from 0.47 µg/l to 7.7 µg/l. In this study, 46.6% of the samples were contaminated with AFB1. The average contamination rate was  $1.1 \pm 0.02$  µg/kg. A total of 6 samples contained more

than 2 µg/kg AFB1, which is within the limit allowed by the Europe Union. As expected, none of them was famous commercial brands.

**Table 1.** Recovery rate obtained from three concentrations

Aflatoxin B1 spiked (µg/kg)	Aflatoxin B1 found (µg/kg)	Recovery (%), n=3
0.6	0.55	91.5 ± 0.16 <sup>a</sup>
1	0.94	94.0 ± 0.23
4	3.64	91.0 ± 0.42

a: Mean ± SD

**Table 2.** List of analyzed tomato pastes

Sample no.	Aflatoxin B1 content (µg/kg) (n=3)
1	0.7 ± 0.035 <sup>b</sup>
2	< LOQ
3	< LOQ
4	0.47 ± 0.016
5	< LOQ
6	< LOQ
7	< LOQ
8	1.6 ± 0.10
9	< LOQ
10	< LOQ
11	< LOQ
12	1.96 ± 0.017
13 <sup>a</sup>	2.33 ± 0.23
14 <sup>a</sup>	7.7 ± 0.29
15 <sup>a</sup>	4.9 ± 0.40
16 <sup>a</sup>	2.9 ± 0.14
17	0.9 ± 0.05
18	< LOQ
19	0.53 ± 0.039
20	< LOQ
21	0.48 ± 0.068
22	< LOQ
23 <sup>a</sup>	2.1 ± 0.06
24	< LOQ
25	< LOQ
26	1.9 ± 0.04
27	< LOQ
28 <sup>a</sup>	3.7 ± 0.10
29	< LOQ
30	< LOQ

a: Means: above the limit allowed by the Europe Union; b: Mean ± SD; LOQ: limit of quantitation

**Table 3.** Compare this study with other researches

Sample	Determination technique	Pretreatment method	LOD	LOQ	Recovery (%)	country	Ref.
Rice	HPLC-FLD	DLLME	0.009 µg/kg	0.03 µg/kg	85.2- 112.0	China	(Lai <i>et al.</i> , 2015)
Noodle	HPLC-FLD	QuEChERS	0.51 µg/kg	1.03 µg/kg	85-109	Malaysia	(Moazami and Jinap, 2009)
Cereals	HPLC-FLD	Validated method	0.0125 ng/g	0.05 ng/g	77-104	Brazil	(Rahmani <i>et al.</i> , 2010)
Black pepper	HPLC-MS/MS	QuEChERS	Not clearly informed	4 µg/kg	102-104	Belgium	(Yogendrarajah <i>et al.</i> , 2013)
Pistachio	HPLC-FLD	(REGULATION, 2006)	0.10 µg/kg	0.11 µg/kg	88.5-89.1	Turkey	(Hepsag <i>et al.</i> , 2014)
Tomato paste	HPLC-FLD	DLLME	0.14 µg/kg	0.44 µg/kg	91-94	Iran	This work

**Table 4.** Comparison of aflatoxin levels with other studies

Samples	No. of samples	% Occurance	Range	Country	Ref.
rice	370	63.5	0.030-20.0 µg/kg	China	(Lai <i>et al.</i> , 2015)
Pistachio	3	66.7	81.6 ng/g	Bahrain	(Musaiger <i>et al.</i> , 2008)
Sunflower seed	50	13	< 2-168 ng/g	Iran	(Beheshti and Asadi, 2013)
Safflower seed	123	85	< 5.4 ng/g		
Pistachio	40	52.5	0.16-122.4 ng/g	Tunisia	(Ghali <i>et al.</i> , 2009)
Sorghum	93	62.0	0.34-52.9 ng/g		
Chicken liver	50	72	0.30-16.36 µg/kg	Iran	(Amirkhizi <i>et al.</i> , 2015)
eggs	150	58	0.30-2.35 µg/kg		
Tomato paste	30	100 (detected) 46.6 (quantified)	< 0.44-7.7 µg/kg	Iran	This work

## Discussion

The advantages of DLLME method include its lower costs due to less solvent consumption, high enrichment factor, and environmental friendliness due to reduced sewage production (Afzali *et al.*, 2012). So far, many methods have been applied for extracting aflatoxin, but the DLLME method is one of the best methods for aflatoxin analysis due to its advantages. **Table 3** shows examples of comparing DLLME with other methods, indicating that DLLME is one of the most appropriate methods for extraction that gives credit to researchers (Chen *et al.*, 2005).

Milk and pistachio are among the most important products which contamination with aflatoxins is frequently reported. Due to the presence of aflatoxin in various foodstuff and

toxicity of AFB1, it is essential to study this mycotoxin in one of the high-consumption foodstuffs in Iran. A study in China showed that 235 out of 370 rice samples were contaminated with AFB1. Moreover, 65 rice samples were positive in terms of contamination with AFB2 (Espinosa-Calderón *et al.*, 2011). Contamination of pistachio nuts with aflatoxins is high in some regions such as the Persian Gulf. A study in Bahrain showed that two of three pistachio samples were contaminated with AFB1 and one of these two samples had a very high contamination level (Afzali *et al.*, 2012). In a study from Iran, AFB1 and AFB2 were detected in 13% and 8% of the sunflower seed samples, respectively. Furthermore, aflatoxins B1, B2, G1, and G2 were detected in 85%, 20%, 16%, and 16% of the

safflower seed samples, respectively. In 10% and 1.6% of the sunflower seeds, the level of AFB1 was above the European Union maximum limit (Campone *et al.*, 2011). In Tunisia, researchers showed that 52.5% of pistachios and 62% of Sorghum included AFB1 positive and the concentrations ranged from < LOQ to 122.4 and 0.34–52.9 ng/g, respectively (Afzali *et al.*, 2012). A study was conducted on 150 eggs and 50 chicken liver samples in Iran. The findings showed that 72% of the chicken liver samples and 87% of the egg samples were contaminated with AFB1 (Kumar *et al.*, 2017). Researchers from the Netherlands and the United Kingdom studied 26 peanuts and found two positive AFB1 samples. **Table 4** contains more information in this regard. These results indicate that the level of aflatoxin in foodstuffs is high throughout the worldwide.

### Conclusions

In this study, an appropriate analytical procedure was successfully developed for analyzing AFB1 in tomato paste samples. As far as we know, no study has ever investigated the presence of aflatoxin in Iran's tomato paste samples. In this study, a rapid and simple analytical procedure was successfully developed for the analysis of aflatoxins in tomato paste samples. Based on the sequential application of solid–liquid extraction and DLLME before the instrumental analysis by HPLC-FLD, this novel method provides a sensitive and accurate determination of analytes.

Based on the findings, AFB1 concentration was above the limit allowed by the Europe Union in the six studied brands. Our aim was to develop an appropriate method to determine aflatoxins in foodstuffs for increasing the humans' health.

### Authors' contributions

Study concept and design: Nemati M and analysis and interpretation of data: Arabkhani P.; drafting of the manuscript: Safavizadeh V, Mojkar M and Shyrina D.

### Conflict of interest

We confirm no known conflicts of interest considering this study.

### Acknowledgements

The authors are grateful to all colleagues that provided support, comments, and inputs. This study was funded by Tabriz University of Medical Sciences.

### References

- Abbrar M, et al.** 2013. Aflatoxins: biosynthesis, occurrence, toxicity, and remedies. *Critical Reviews in Food Science and Nutrition*. **53 (8)**: 862-874.
- Afzali D, Ghanbarian M, Mostafavi A, Shamspur T & Ghaseminezhad S** 2012. A novel method for high preconcentration of ultra trace amounts of B1, B2, G1 and G2 aflatoxins in edible oils by dispersive liquid–liquid microextraction after immunoaffinity column clean-up. *Journal of Chromatography A*. **1247**: 35-41.
- Alshannaq A & Yu J-H** 2017. Occurrence, toxicity, and analysis of major mycotoxins in food. *International Journal of Environmental Research and Public Health*. **14 (6)**: 632.
- Amirkhizi B, Arefhosseini SR, Ansarin M & Nemati M** 2015. Aflatoxin B1 in eggs and chicken livers by dispersive liquid–liquid microextraction and HPLC. *Food Additives & Contaminants: Part B*. **8 (4)**: 245-249.
- Authority EFS** 2007. Opinion of the scientific panel on contaminants in the food chain [CONTAM] related to the potential increase of consumer health risk by a possible increase of the existing maximum levels for aflatoxins in almonds, hazelnuts and pistachios and derived products. *EFSA Journal*. **5 (3)**: 446.
- Beheshti HR & Asadi M** 2013. Aflatoxins in sunflower and safflower seeds from Iran. *Food Additives and Contaminants: Part B*. **6 (1)**: 68-71.
- Berijani S, Assadi Y, Anbia M, Hosseini M-RM & Aghaee E** 2006. Dispersive liquid–liquid microextraction combined with gas chromatography-flame photometric detection: very simple, rapid and sensitive method for the determination of organophosphorus pesticides in

- water. *Journal of Chromatography A*. **1123** (1): 1-9.
- Campone L, Piccinelli AL, Celano R & Rastrelli L** 2011. Application of dispersive liquid-liquid microextraction for the determination of aflatoxins B1, B2, G1 and G2 in cereal products. *Journal of Chromatography A*. **1218** (42): 7648-7654.
- Chen J, Zhang X, Yang M & Jin Y** 2005. Recent advances in the determination methods of aflatoxins. *Journal of Chinese Materia Medica*. **30** (24): 1890-1894.
- Espinosa-Calderón A, et al.** 2011. Methods for detection and quantification of aflatoxins. Aflatoxins: detection, measurement and control. InTech: New York.
- Flores-Flores ME, Lizarraga E, de Cerain AL & González-Peñas E** 2015. Presence of mycotoxins in animal milk: A review. *Food Control*. **53**: 163-176.
- Fontana AR, Prendes LP, Morata VI & Bottini R** 2016. High-throughput modified QuEChERS method for the determination of the mycotoxin tenuazonic acid in wine grapes. *RSC Advances*. **6** (98): 95670-95679.
- Ghali R, et al.** 2009. Simultaneous HPLC determination of aflatoxins B1, B2, G1 and G2 in Tunisian sorghum and pistachios. *Journal of Food Composition and Analysis*. **22** (7-8): 751-755.
- Hepsag F, Golge O & Kabak B** 2014. Quantitation of aflatoxins in pistachios and groundnuts using HPLC-FLD method. *Food Control*. **38**: 75-81.
- Kilicel F, Karapinar HS & Cimen A** 2017. Quantitation of Aflatoxins in Food Materials Using HPLC-FLD Method. *Science Journal of Analytical Chemistry*. **5** (6): 90.
- Kumar P, Mahato DK, Kamle M, Mohanta TK & Kang SG** 2017. Aflatoxins: a global concern for food safety, human health and their management. *Frontiers in Microbiology*. **7**: 2170.
- Lai X, Liu R, Ruan C, Zhang H & Liu C** 2015. Occurrence of aflatoxins and ochratoxin A in rice samples from six provinces in China. *Food Control*. **50**: 401-404.
- Moazami E & Jinap S** 2009. Natural occurrence of deoxynivalenol (DON) in wheat based noodles consumed in Malaysia. *Microchemical Journal*. **93** (1): 25-28.
- Musaiger AO, Al-Jedah JH & D'souza R** 2008. Occurrence of contaminants in foods commonly consumed in Bahrain. *Food Control*. **19** (9): 854-861.
- Rahmani A, Jinap S & Soleimany F** 2010. Validation of the procedure for the simultaneous determination of aflatoxins ochratoxin A and zearalenone in cereals using HPLC-FLD. *Food Additives & Contaminants: Part A*. **27** (12): 1683-1693.
- Rawat S** 2015. Food Spoilage: Microorganisms and their prevention. *Asian Journal of Plant Science and Research*. **5** (4): 47-56.
- REGULATION HAT** 2006. Commission Regulation (EC) No 1259/2007. *Journal of Investig Allergol ClinIcal Immunology*. **16**: 136-137.
- Rezaee M, et al.** 2006. Determination of organic compounds in water using dispersive liquid-liquid microextraction. *Journal of Chromatography A*. **1116** (1-2): 1-9.
- Yogendrarajah P, Van Poucke C, De Meulenaer B & De Saeger S** 2013. Development and validation of a QuEChERS based liquid chromatography tandem mass spectrometry method for the determination of multiple mycotoxins in spices. *Journal of Chromatography A*. **1297**: 1-11.
- Yu J, et al.** 2004. Genomics of economically significant *Aspergillus* and *Fusarium* species. *Applied Mycology and Biotechnology*. **4**: 249-283.