Journal of Nutrition and Food Security
Shahid Sadoughi University of Medical Sciences
School of Public Health
Department of Nutrition
Nutrition & Food Security Research Center

eISSN: 2476-7425 pISSN: 2476-7417 JNFS 2017; 2(2): 135-137 Website: jnfs.ssu.ac.ir

Vitamin D Deficiency: A Neglected Truth in Iran

Nahid Ramezani-Jolfaie; MSc [1,2] & Amin Salehi-Abargouei; PhD[1,2*]

[1] *Nutrition and Food Security Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.*
[2] *Department of Nutrition, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.*

ARTICLE INFO

EDITORIAL ARTICLE

Article history:
Received: 17 Nov2016
Revised: 21 Dec 2016
Accepted: 21 Jan 2017

Corresponding author:
Department of Nutrition, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
abargouei@ssu.ac.ir

Postal code: 8916877391
Tel: +98-35-31492229

Vitamin D (vit D) is a fat-soluble steroid which includes two forms: vit D2 (ergocalciferol) and vit D3 (cholecalciferol). The former is provided from plant and the latter is taken from animal sources (Shao *et al.*, 2012). Vit D is obtained from natural synthesis in the skin and also from some dietary sources such as oily fish, eggs, fortified products (dairy, cereals, soy), and oral supplements (Jolfaie *et al.*, 2016, Kennel *et al.*, 2010). Finally, it is activated biologically during hydroxylation process in the liver and kidneys (Shao *et al.*, 2012).

Vit D has essential physiological functions for maintaining mineral homeostasis (Chung *et al.*, 2011). Indeed, vit D deficiency can contribute to several musculoskeletal disorders, such as rickets in children, osteomalacia, osteopenia, osteoporosis, and fractures in adults (Moher *et al.*, 2001). In addition, the association between vit D deficiency and many diseases including cancers, infections, autoimmune diseases, diabetes mellitus, and cardiovascular diseases have been studied (Hilger *et al.*, 2014, Norman, 2008). The normal level of 25(OH)D (the dominant vit D metabolite in circulation representing vit D status) is considered 20 ng/ml (50 nmol/L) and less than this value is classified into the deficiency category (Christodoulou *et al.*, 2013, DeLuca, 2004). The most important factors in deficiency of vit D are decreased sun exposure, insufficient vit D intake, and increase of age (Gaugris *et al.*, 2005). Moreover, high risk of vit D deficiency has been observed in special groups and conditions such as children, pregnant and postmenopausal women, as well as elderly people (Shao *et al.*, 2012); diseases associated with fat mal-absorption (e.g., Crohn's disease) (Kennel *et al.*, 2010); and disturbing conditions of parathyroid, liver, or kidney functions (Kathleen Mahan *et al.*, 2012).

This paper should be cited as: Ramezani-Jolfaie N, Salehi-Abargouei A. ***Vitamin D Deficiency: A Neglected Truth in Iran.*** *Journal of Nutrition and Food Security (JNFS)*, 2017; 2 (2): 135-137.

[Downloaded from jnfs.ssu.ac.ir on 2026-11-28]
[DOR: 20.1001.1.24767417.2017.2.2.2.8]

On the basis of current reports, there are approximately one billion people with vit D deficiency worldwide (Khadilkar and Khadilkar, 2013). In Iran, despite the fact that it is a sunny country, the high prevalence of vit D deficiency is a growing concern (Heshmat *et al.*, 2008). According to a population-based study in our country, the rate of vit D deficiency among women and men has been estimated to be about 75.1 % and 72.1%, respectively (Moradzadeh *et al.*, 2008). Furthermore, Yazd as a province in center of Iran is located in desert areas (in the latitude of 31.5^0) and is thus mostly sunny in the year, has prevalent vit D deficiency. Some studies have indicated that 60% of girl students and 78 % of medical staffs of Shahid Sadughi hospital had deficiency (Mehrdad *et al.*, 2009, Shakiba and Rafiei, 2009). As mentioned earlier, several studies indicated the association between vit D status and chronic diseases; however, in Iran the studies did not replicate these results. This might be cause by the fact that all participants had vit D deficiency and there was low variation between them in serum vit D levels. Frequent use of topical sunscreen by women and lack of vit D fortified foods might be some other reasons for high prevalence of vit D deficiency in Iran despite its sunny climate (Heshmat *et al.*, 2008).

Some recommendations have been suggested that can help to overcome this situation. Measurement of serum 25(OH)D levels regularly to identify the necessity of vit D supplementation. Daily sunscreen-free exposure to the sun (10-15 minutes) to provide vit D. Consumption of foods that naturally contain vit D (salmon, sardines mackerel, and cod liver oil). Consumption of fortified foods with vit D3 (e.g., fortified milk, orange juice, formulas, yogurts [100 IU/8 oz], cheeses [100 IU/3 oz], butter [56 IU/3.5 oz], margarine [429 IU/3.5 oz], and breakfast cereals [100 IU/serving]). Using vit D supplementation to treat vit D deficiency and maintenance therapy. In vit D supplementation mentioned above, the recommended supplementation doses were as the following: infants aged 0–1 year [2000 IU/d, 400-1000 IU/d], children aged 1–18 years [2000 IU/d, 600-1000 IU/d], adults [50,000 IU/week for 8 weeks, 1500–2000 IU/d]) (Holick *et al.*, 2011). Moreover, the results of meta-analysis studies indicated that 700-800 IU/d vit D supplementation can reduce the risk of fractures and bone loss in elderly people (Bischoff-Ferrari *et al.*, 2005, Tang *et al.*, 2007). The results of another meta-analysis also revealed that serum vit D levels are inversely associated with all-cause and cause-specific mortality which impose significant economic burden on the healthcare system (Chowdhury *et al.*, 2014, Schöttker *et al.*, 2014). According to findings of the study conducted on Canadian population, the economic burden could be decreased to 7.3% (4.0-10.5%) or $14.4 billion ($8.0–$20.1 billion) if the mean serum vit D concentration increases to 105 nmol/L (Grant *et al.*, 2010). Therefore, regarding the importance of vit D effects on body health and subsequently health care system, it is necessary to pay special attention to vit D status in all age groups.

Authors' Contributions

Ramezani-Jolfaie N wrote the manuscript. Salehi-Abargouei A edited the manuscript. Both authors approved the content of the manuscript, and agreed for all aspects of the work.

References

Bischoff-Ferrari HA, et al. 2005. Fracture prevention with vitamin D supplementation: a meta-analysis of randomized controlled trials. *Journal of the American medical association.* **293 (18)**: 2257-2264.

Chowdhury R, et al. 2014. Vitamin D and risk of cause specific death: systematic review and meta-analysis of observational cohort and randomised intervention studies. *British medical journal* **348**: g1903.

[Downloaded from jnfs.ssu.ac.ir on 2026-11-28]

[DOR: 20.1001.1.24767417.2017.2.2.2.8]

Christodoulou S, Goula T, Ververidis A & Drosos G 2013. Vitamin D and bone disease. *BioMed research international.* **2013**: 396541.

Chung M, Lee J, Terasawa T, Lau J & Trikalinos TA 2011. Vitamin D with or without calcium supplementation for prevention of cancer and fractures: an updated meta-analysis for the US Preventive Services Task Force. *Annals of internal medicine.* **155 (12)**: 827-838.

DeLuca HF 2004. Overview of general physiologic features and functions of vitamin D. *The American journal of clinical nutrition.* **80 (6)**: 1689S-1696S.

Gaugris S, et al. 2005. Vitamin D inadequacy among post-menopausal women: a systematic review. *Monthly journal of the association of physicians.* **98 (9)**: 667-676.

Grant WB, Schwalfenberg GK, Genuis SJ & Whiting SJ 2010. An estimate of the economic burden and premature deaths due to vitamin D deficiency in Canada. *Molecular nutrition & food research.* **54 (8)**: 1172-1181.

Heshmat R, et al. 2008. Vitamin D deficiency in Iran: A multi-center study among different urban areas. *Iran journal public health.* **37 (suppl)**.

Hilger J, et al. 2014. A systematic review of vitamin D status in populations worldwide. *British journal of nutrition.* **111 (01)**: 23-45.

Holick MF, et al. 2011. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *The Journal of clinical endocrinology & metabolism.* **96 (7)**: 1911-1930.

Jolfaie NR, Rouhani MH, Onvani S & Azadbakht L 2016. The association between Vitamin D and health outcomes in women: A review on the related evidence. *Journal of research in medical sciences: The official journal of Isfahan University of medical sciences.* **21**: 76.

Kathleen Mahan L, Escott-Stump S, Raymond Janice L & Krause MV 2012. Krause's Food & the Nutrition Care Process. Elsevier health sciences.

Kennel KA, Drake MT & Hurley DL 2010. Vitamin D deficiency in adults: when to test and how to treat. In *mayo clinic proceedings*. **85(8)**: 752-758. Elsevier.

Khadilkar VV & Khadilkar AV 2013. Use of vitamin D in various disorders. *The Indian journal of pediatrics.* **80 (3)**: 215-218.

Mehrdad S, Zahra N, Hassan LM & Ahmad S 2009. Prevalence of vitamin D deficiency among female students in secondary guidance school in Yazd City. *Acta medica Iranica.* **47 (3)**: 209-214.

Moher D, Schulz KF, Altman D & Group C 2001. The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomized trials. *Journal of the American medical association.* **285 (15)**: 1987-1991.

Moradzadeh K, et al. 2008. Normative values of vitamin D among Iranian population: a population based study. *International journal of osteoporosis & metabolic disorders.* **1 (1)**: 8-15.

Norman AW 2008. A vitamin D nutritional cornucopia: new insights concerning the serum 25-hydroxyvitamin D status of the US population. *The American journal of clinical nutrition.* **88 (6)**: 1455-1456.

Schöttker B, et al. 2014. Vitamin D and mortality: meta-analysis of individual participant data from a large consortium of cohort studies from Europe and the United States. *British medical journal* **348**: g3656.

Shakiba M & Rafiei P 2009. Prevalence of vitamin D deficiency among medical staff in Shahid Sadoughi Hospital in Yazd, Iran. *British medical journal.* **7 (25)**: 22-30.

Shao T, Klein P & Grossbard ML 2012. Vitamin D and breast cancer. *The oncologist.* **17 (1)**: 36-45.

Tang BM, Eslick GD, Nowson C, Smith C & Bensoussan A 2007. Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. *The lancet.* **370 (9588)**: 657-666.

[Downloaded from jnfs.ssu.ac.ir on 2026-11-28]

[DOR: 20.1001.1.24767417.2017.2.2.2.8]

[DOR: 20.1001.1.24767417.2017.2.2.2.8] [Downloaded from jnfs.ssu.ac.ir on 2026-11-28]