



Journal of Nutrition and Food Security

Shahid Sadoughi University of Medical Sciences
School of Public Health
Department of Nutrition
Nutrition & Food Security Research Center

eISSN: 2476-7425

pISSN: 2476-7417

JNFS 2024; 9(1): 132-143

Website: jnfs.ssu.ac.ir

Pesticide Residue in Iranian Fruits and Vegetables: A Systematic Review

Fatemeh Toorang; PhD^{1,3}, Bahareh Sasanfar ; PhD*^{2,4,6}, Hamed Pouraram; PhD¹ & Soheyl Eskandari; PhD⁷

¹ Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran; ² Cancer Research Center, Cancer Institute of Iran, Tehran University of Medical Sciences, Tehran, I.R. Iran; ³ Departments of Medical and Surgical Sciences, University of Bologna, Italy; ⁴ Department of Nutrition, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran; ⁵ Research Center for Food Hygiene and Safety, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran; ⁶ Student Research Committee, Shahid Sadoughi University of Medical Sciences, Yazd, Iran; ⁷ Food and Drug Laboratory Research Center, Food and Drug Administration, Iran Ministry of Health and Medical Education, Tehran, Iran.

ARTICLE INFO

SYSTEMATIC REVIEW

Article history:

Received: 6 Jul 2022

Revised: 29 Jul 2022

Accepted: 28 Aug 2022

*Corresponding author:

bsasanfar@yahoo.com

Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran.

Postal code: 13145-158

Tel: +98 21 66581638

ABSTRACT

Background: There is a growing concern about the health-threatening effects of pesticide residues in fruits and vegetables worldwide. This study systematically reviewed the published data on pesticide residues in Iranian fruits and vegetables to clarify the gap in this issue. **Method:** The authors systematically searched PubMed, Google Scholar, Scopus, SID, and Iran Medex to find published studies on pesticide residues in Iranian foods without time and language restrictions. The title and abstract of all articles were evaluated after removing duplicate articles (2289 articles) by two independent reviewers. Finally, 24 articles were found that reported pesticide residues in fruits and vegetables. There was a great variation in measurement methods and pesticides reported across studies, which precluded meta-analysis. Therefore, a summary of the included studies was only reported. **Results:** Twenty-four studies reporting pesticide residues in Iranian fruits and vegetables were included. The percentage of Iranian fruits and vegetables contaminated with pesticides exceeding the maximum residue limit (MRL) was less than 10% in most studies. Contaminated samples were collected mainly from cultivated areas such as fields, orchards, or greenhouses. **Conclusion:** Pesticide residues in food have not been systematically reported in Iran. It was found that only limited articles were published by academic societies on this issue. Considering the current scenario, there is an urgent need to facilitate reliable and continuous measurements of toxic residues in Iranian food.

Keywords: Pesticide; Fruit; Vegetable; Systematic review; Iran

Introduction

Fruits and vegetables are the main components of a healthy diet recommended by international guidelines to improve general health and reduce the risk of several diseases

(World Health Organization, 2019). Many studies have identified plant food as food rich in various micronutrients and phytochemicals that are essential for health and prevention of common

This paper should be cited as: Toorang F, Sasanfar B, Pouraram H, Eskandari S. *Pesticide Residue in Iranian Fruits and Vegetables: A Systematic Review. Journal of Nutrition and Food Security (JNFS), 2024; 9 (1): 132-143.*

diseases, such as cancer and cardiovascular diseases (Mayne *et al.*, 2016). Moreover, replacing animal food with fruits and vegetables can protect our planet (Vaidyanathan, 2021).

There is a growing concern about the health-threatening effects of pesticide residues in fruits and vegetables worldwide. It is clear that pesticides harm the environment and every living being, including humans. Studies have shown that exposure to pesticides may lead to chronic and short-term health problems in humans. The most common short-term side effects are nausea, vomiting, and headache (Kim *et al.*, 2017). Long-term effects that are not easily detected and can be more dangerous include behavioral changes, several cancers, and disruption of the endocrine and reproductive systems (Alavanja *et al.*, 2004, Asghar *et al.*, 2016, Kim *et al.*, 2017).

On the other hand, most farmers and policymakers believe that pesticides are important components in improving the food security of countries and their use is an inevitable factor to meet the global demand for sufficient and affordable food (Popp *et al.*, 2013). Considering these two arguments, some international organizations, including the World Health Organization (WHO), have established a standard protocol for the use of pesticides, and the maximum residue limit (MRL) for pesticides in food, which is expressed in milligrams per kilogram of food, and if a food has a higher limit, it will be considered unhealthy (Herrman, 1993).

Developing countries such as Iran have been accused of excessive use of toxins, which is generally due to the identification of some high levels of residual toxins in export food baskets. This issue can damage their food image all over the world (Donkor *et al.*, 2016). Moreover, the level of pesticide residues in domestic food is likely to be similar to that of exports, which could affect national health (Carvalho, 2006). These problems may be due to the lack of awareness of food producers, including farmers, as well as ineffective national laws and permits (Carvalho, 2006).

Due to the mentioned danger of pesticides for

human health and wildlife, the use of pesticides is under continuous monitoring. Although there have been studies on pesticide residues in Iran, there is no comprehensive information about it. Therefore, this study systematically reviewed the published data on pesticide residues in Iranian fruits and vegetables to clarify the gap in this issue.

Materials and Methods

Data sources: PubMed, Google Scholar, Scopus, SID, and Iran Medex were systematically searched to find published studies on pesticide residues in Iranian foods without time and language restrictions. Different search strategies were used in different databases (**Table 1**) using these keywords:

(Pesticide or herbicide or insecticide) and Iran

Study selection: The title and abstract of all articles were evaluated after removing duplicate articles (2289 articles) by two independent reviewers and resolved disagreements through discussion. The title and abstract of the articles found in Google Scholar were read and 26 articles that met the study criteria were entered into the Endnote file. Reference lists of included studies were reviewed to identify articles not captured by the authors' search.

This study aimed to systematically review all the studies that have reported the residual toxins in Iranian food. After reading the abstracts of the articles, the researchers included 30 studies on fruits and vegetables, 10 on fish, 2 on honey, and 27 on other food types. Considering the comprehensiveness of the data related to fruits and vegetables, as well as the Iranian society's fear of agricultural pesticide residues in fruits and vegetables, only the results related to the pesticide residues in fruits and vegetables were reported. After reading the full text of 30 articles on fruits and vegetables, 24 articles met the study criteria. One study related to laboratory-grown vegetables, three studies on standardized methods for measuring toxins, one study on fungal contamination, and one study on phthalate contamination were excluded.

Data extraction: Data from 24 studies related to

fruits and vegetables were transferred to Excel tables. The data were summarized based on a table prepared by two reviewers, which included data on the type of vegetables or fruits, pesticides identified, first author and year of publication, geographical location, mean concentration, percentage of contaminated samples, and the place of sample collection (farm, market, or garden). There was a great variation in measurement methods and pesticides reported across studies, which precluded meta-analysis. Therefore, the authors only reported summary findings of the included studies.

Results

First, 2464 articles were found from international electronic databases, 205 of which were duplicate articles, and also 30 articles were found from Persian databases. Therefore, the title and abstract of 2289 articles were reviewed to find studies that investigated pesticide residues in Iranian food. Finally, 24 articles were found that reported pesticide residues in fruits and vegetables (24 articles), wheat (1 article), rice (1 article), tea (1 article), sugar (1 article), eggs (1 article), wild duck (1 article), fish (5 articles), and honey (1 article). Since there was limited information on food groups other than fruits and vegetables, this report was limited to fruits and vegetables.

Most studies measured pesticide residues in samples obtained from gardens, greenhouses, and fields (Behbahaninia, 2007, Ganjeizadeh Rohani *et al.*, 2014, Hagian Shahri *et al.*, 2014, Khak *et al.*, 2016, Khaniki *et al.*, 2011, Leili *et al.*, 2016, Pirsahab *et al.*, 2017, Sobhanardakani *et al.*, 2016). Two studies measured pesticide residues in samples obtained simultaneously from market and cultivated areas (Sobhanardakani *et al.*, 2014, Sobhanardakani *et al.*, 2016). Jahanmard *et al.* examined tomatoes from a salad factory. This was the only study that tested the residue after processing (Jahanmard *et al.*, 2016). Finally, some studies reported residuals in market samples (Akhlaghi *et al.*, 2013, askari *et al.*, 2014, Hadian and Azizi, 2008). The residual level in the cultivation areas was generally higher than the market.

Most of the studies investigated pesticide residues in cucumber and tomato, as 9 studies were conducted on cucumber (Ardakani *et al.*, 2012, Behbahaninia, 2007, Farshad, 2001, Ganjeizadeh Rohani *et al.*, 2014, Hadian *et al.*, 2006, Hagian Shahri *et al.*, 2014, Khaniki *et al.*, 2011, Leili *et al.*, 2016, Shokrzadeh *et al.*, 2013) and 9 studies on tomato (Ardakani *et al.*, 2012, Bayat *et al.*, 2015, Hadian *et al.*, 2006, Hagian Shahri *et al.*, 2014, Jafari *et al.*, 2012, Jahanmard *et al.*, 2016, Khak *et al.*, 2016, Khaniki *et al.*, 2011, Mohammadi and Imani, 2012). Other studied commodities included apples (Akhlaghi *et al.*, 2013, Hagian Shahri *et al.*, 2014, Mina and Maryam, 2012, Pirsahab *et al.*, 2017), melons (Akhlaghi *et al.*, 2013, Hadian and Azizi, 2008), grapes (Hagian Shahri *et al.*, 2014), mushrooms (Sobhanardakani *et al.*, 2014), zucchini (Sobhanardakani *et al.*, 2016), strawberry (Golepoor *et al.*, 2014) cherry (Akhlaghi *et al.*, 2013, askari *et al.*, 2014), and watermelons (Akhlaghi *et al.*, 2013). In 24 studies, 36 pesticide residues were reported in different commodities. They included abamectin, diazinon, chlorpyrifos, ethion, imidacloprid, cypermethrin, permethrin, indoxacarb, mancozeb, chlorothalonil, iprodione, thiophanate methyl, carbendazim, gelsam, pyrethroids, dischlorompatiraphan, tepe, dichloropatyrphan, tepe, dichloropatylate, carben, dischlorofandoparithion, carblo, endosulfan I, endosulfan II, endosulfan sulfate, oxymethon, methyl, dichlorvos, metalaxyl, fenpropathrin, fenpropathrin, malate, fenitrothion, oxymethon methyl, P metrozine, fesalone, and fenvalerate.

Unfortunately, studies mostly reported only pesticide concentrations, and most of them did not report the percentage of commodities that were contaminated above the MRL. In most of the studies reporting the percentage exceeding the MRL, it was less than ten percent of the commodity. However, two studies reported a high percentage of heavily contaminated commodities. Ganjeizadeh reported that 53% of Kerman greenhouse cucumbers were contaminated with diazinon more than the MRL, and this percentage was 78.33% for methyl oxymethon (Ganjeizadeh Rohani *et al.*, 2014). Diazinon contamination was

reported in all cucumbers and tomato greenhouses of Chaharmahal and Bakhtiari provinces exceeding the MRL (Khaniki *et al.*, 2011). It should be noted

that some studies investigated the residues of banned pesticides such as carbamil and fortunately these products were free of those pesticides.

Table 1. Search strategies in different databases.

Database	Searching strategy	Number of found articles	Date of search
Scopus	(Pesticide OR herbicide OR insecticide [title/abstract]) AND Iran (affiliation)	2066	2018 July 8
PubMed	(Pesticide OR herbicide OR insecticide [title/abstract]) AND Iran (affiliation)	389	2018 July 8
Google Scholar	(Pesticide herbicide insecticide)+ Iran	17500	2018 July 8
Sid	Pesticide or herbicide (in Persian)	25	2018 July 8
Iran medex	Pesticide or herbicide (in Persian)	5	2018 July 8




Figure 1. Flow diagram of study screening.

Table 2. Pesticides detected in fruits and vegetables in Iran.

Author	Sampling site	Sample size	Detecting technique	Geographic location	Commodity	Mean or range of concentrations(ppm) of Detected pesticides
(Pirsaheb <i>et al.</i> , 2017)	Gardens	500g of golden or red apples from 50 garden	DLLME-SFO coupled with HPLC-UV	Mahabad	Golden apple Red apple	Abamectin (0), Diazinon (10.289), Chlorpyrifos (9.51) Abamectin (0), Diazinon (8.867), Chlorpyrifos (8.047)
(Leili <i>et al.</i> , 2016)	Greenhouses	1000mg from 10 greenhouses	QuEChERS Followed by GC-MS	Hamadan	Cucumber (One day after pesticide application)	Ethion (0.867-0.975), Imidacloprid (1.13-1.207)
(Sobhanardakani <i>et al.</i> , 2016)	Greenhouses and markets		Spectrophotometric	Hamadan	Zucchini	Diazinon (0.093-0.159)
(Khak <i>et al.</i> , 2016)	Farms	37 samples from every farm		Jam, Dashtestan, Dashti, Deir, and Kangan (mean of all cities are reported)	Tomato	Cypermethrin (0.071±0.069), Permethrin (0.272±0.19), Indoxacarb (0.03±0.017), Mancozeb (0.035±0.019), Chlorothalonil (0.15±0.238), Iprodione (0.004±0.003), Thiophanate Methyl (0.11±0.183), Carbendazim golsam (0.107±0.156), Abamectin (0)
(Jahanmard <i>et al.</i> , 2016)	Salad production plant	22 samples	QuEChERS Followed by GC-MS	Isfahan	Tomato	Pyrethroid, Diazinone (107.67-579.81), Chlorpyrifos (144.92-254.84), Primicarb (free), Dischlorvos(free), Carbaryl(free), Malathion(free), Brompropilate (free), Propargit (free), Tetradifone(free), Posalone(free), Iprodion(free), Endosulfane(free)
(Sobhanardakani <i>et al.</i> , 2014)	Greenhouses and markets	10 samples	Spectrophotometry	Hamadan	Mushroom	Diazinon (0.04-0.166)
(Hagian Shahri <i>et al.</i> , 2014)	Cultivation regions	6 samples of cucumber	QuEChERS Followed by GC-MS	Mashhad, Neyshaboor and Sabzevar	Cucumber	Malathion (5.09), Oxydemethon (3.33), Methyl (0.18), Diazinon (0.43), Dichlorvos (2.38), Metalaxyl(0.41), Fenpropathrin(6.32),
		8 samples of tomato		Mashhad, Chenaran, Fariman, Neyshaboor and Ghoochan	Tomato	Fanpropatrin (7.65), Azinphos methyl (0.16), Diazinon (0.235), Phosalone (0.1),
		8samples of cherries		Mashhad and Neyshaboor	Cherry	Diazinon (11.2), Malation (0.79), Fenpropidin (1.57),
		10 samples of grapes		Hagian-shahri M.	Grape	Ethion (2.4), Malathion (12.46), Diazinon (0.48)
		7 samples of apples		Mashhad, Chenaran, Ghoochan and Neyshaboor	Apple	Ethion (1.59), Fenitrothion (0.06)

(Ganjeizadeh Rohani et al., 2014)	Greenhouses	60 samples	spectrophotometry	Kerman	cucumber	Diazinon (0.582), Oxydemeton methyl (1.91)
(Akhlaghi et al., 2013)	Local markets and villages	75	QuEChERS Followed by GC-MS	A change village	Apple	0.17±0.04
					Grape	0.41±0.15
					Melon	0.27±0.04
					Watermelon	0.12±0.03
					Cherry	0.18±0.04
					Apple	0.25±0.08
				Tabas village	Grape	0.22±0.06
					Melon	0.27±0.06
					Watermelon	0.20±0.06
					Cherry	0.11±0.06
					Apple	0.14±0.04
					Grape	0.35±0.12
				Barghamad village	Melon	0.84±0.19
					Watermelon	0.18±0.06
					Cherry	0.30±0.18
					Apple	0.26±0.11
					Grape	0.30±0.14
					Melon	0.55±0.14
Sabzevar local market	Watermelon	0.23±0.08				
	Cherry	0.88±0.21				
	Apple	0.24±0.07				
	Grape	0.34±0.15				
Neyshabur local market	Melon	0.26±0.09				
	Watermelon	0.17±0.07				
	Cherry	0.12±0.02				
(Khaniki et al., 2011)	Greenhouses	60 samples	HPLC	Chaharmahal and Bakhtiari province	Cucumber	Oxydimeton methyl 0.23±0.17
					Tomato	1.61±0.8

(askari <i>et al.</i> , 2014)	Wholesale markets	40 samples		Tehran (From farms in Lavasan, Shahriar, Ghazvin, Mashhad, and Orumieh)	Cherry	Diazinon Mashhad: 0.3Lavasan: 0.29Diazinon in Samples from other cities were undetectable
(Behbahaninia, 2007)	Farms	12 samples	Gas chromatography	Damavand	Cucumber	Tetradifone (0-0.92), Pymetrozine (0.0214-2.67), Deltamethrin (0-0.55)
(Hadian and Azizi, 2006)	Wholesale markets	10 samples of cucumber 10 samples of tomato	GC/ITMS To detect 117 pesticides	Tehran market (from Poldokhtar farms)	Cucumber	Endosulfane I (0.032±0.0049), Endosulfane II (0.03±0.0045), Endosulfane sulphate (0.04±0.0032)
				Tehran market (from Khorramabad farms)	Cucumber	Chlorpyrifos (0.028±0.0037)
				Tehran market (from Ghazvine farms)	Tomato	Phosalone (0.045±0.0084)
				Tehran market (from Varamin farms)	Tomato	Fenvalerate (0.05),
(Golepoor <i>et al.</i> , 2014)	Farms	3	QUECHERS GC-MS	Tonekabon	Strawberry	Dursban (ND), Diazinon (ND), Ethion (ND), Malathion (ND)
		120		Bahmanmir-Babolsar		Dursban (3.47±0.52), Diazinon (9.10±1.49), Ethion (1.35±0.06), Malathion (7.99±0.93)
		5		Amol		Dursban (ND), Diazinon (ND), Ethion (ND), Malathion (ND),
		15		Babol		Dursban (8.82±0.82), Diazinon (ND), Ethion (ND), Malathion (ND)
		28		Jooybar		Dursban (9.99 ±1.65), Diazinon (1.84±0.29), Ethion (12.15 ±1.15), Malathion (ND)
		10		Kiakola		Dursban (ND), Diazinon (6.33±1.33), Ethion (6.23±0.23), Malathion (ND)
10	Behshahr	Dursban(ND), Diazinon(219.68±32.65), Ethion(ND), Malathion (0.56±0.04),				
20	Sari	Ethion (ND), Malathion (ND)				
(Mohammadi and Imani, 2012)	Wholesale markets	2000mg in 25 samples from 10 markets	HPLC GC/NPD GC/MS	Karaj	Tomato	Chlorpyrifos (0.2), Deltamethrin (0.09)
(Jafari <i>et al.</i> , 2012)	Central fruit and vegetables market	40 greenhouses,40 greenhouse from different cities	HPLC with UV detection	Tehran	Tomato	Dithiocarbamate (0.14)
(Bayat <i>et al.</i> , 2015)	The market of Mashhad City	4 samples	HPLC	Hormozgan	Tomato	Diazinon 0.20±0.01
				Khuzestan		0.36±0.01

				Shiraz		0.46±0.01
				Neyshaboor		0.54±0.01
				Chenaran		0.57±0.02
				Mashhad		0.64±0.01
(Mina and Maryam, 2012)	Garden	972 samples		Damavand	Red apple	Diazinon (0.70±0.36), Chlorpyrifos (1.35±0.82),
					Golden apple	Diazinon (0.65±0.17), Chlorpyrifos (1.09±0.21),
(Hadian et al., 2006)	Central fruit and vegetables market in Tehran	10	HPGPC CC/ITMS	Saveh	Cantaloupe	Endosulfan II (0), Endosulfan sulfate (0.06)
		10		Dezful	Watermelon	Endosulfan II (ND), Endosulfan sulfate (ND)
		10		Ahvaz		Endosulfan II(ND), Endosulfan sulfate (ND)
		10		Varamin	Melon	Endosulfan II(ND), Endosulfan sulfate (ND)
				Torbatejam		Endosulfan II(ND), Endosulfan sulfate (0.02)
(Hadian and Azizi, 2008)	Central fruit and vegetables market in Tehran	10 samples of cucumber	GC/ITMS	Poldokhtar	Cucumber	Endosulfan I (0.032±0.005), Endosulfan II (0.03±0.004)
		10 samples of tomato		Khorramabad		Endosulfan sulphate (0.04±0.003), Chlorpyrifos (0.028±0.004)
(Shokrzadeh et al., 2013)	Garden	100	GC/MS	Varamin	Tomato	Phosalone (0.45±0.008) Fenvalerate (0.05)
		8 samples		Sari	Orange	Diazinon (0.4)
		20 samples		Neka (8 samples)		Benomyl: bush (0.037±0.002), tree (0.043±0.003); Mancozeb: bush (0.029±0.004), tree (0.030±0.03)
(Shokerzadeh et al., 2006)	Farm	16	GC	Sari (20 samples)		Benomyl: bush (0.21±0.001), tree (0.039±0.003); Mancozeb: bush (0.033±0.002), tree (0.039±0.004)
		12		Jooybar (16 samples)	Cucumber	Benomyl: bush (0.038±0.004), tree (0.041±0.002); Mancozeb: bush (0.046±0.002), tree (0.035±0.003)
		4		Ghaemshahr (12 samples)		Benomyl: bush (0.028±0.003), tree (0.032±0.001); Mancozeb: bush (0.030±0.001), tree (0.021±0.002)
				Babol (4 samples)		Benomyl: bush (0.024±0.01), tree (0.042±0.003); Mancozeb: bush (0.031±0.001), tree (0.034±0.004)

		8		Babolsar(8 samples)		Benomyl: bush (0.051±0.004), tree (0.026±0.002); Mancozeb: bush (0.035±0.002), tree (0.048±0.005)
		72		Total (mazandarn province) (72 samples)		Benomyl: bush (0.032±0.002), tree (0.036±0.002); Mancozeb: bush (0.035±0.002), tree (0.036±0.003)
(Ardakani <i>et al.</i> , 2012)	Farm	106 samples of cucumber and 48 samples of tomato	GC(ECD/NPD)	Gachsaran	Cucumber	Endosulfan: α isomer (0.352), β isomer (0.443); Diazinon (0.462)
					Tomato	Endosulfan: α isomer (0.325), β isomer (0.284); Diazinon (0.504)
				Kohgyloveh	Cucumber	Endosulfan: α isomer (0.167), β isomer (0.271); Diazinon (0.205)
					Tomato	Endosulfan: α isomer (0.130), β isomer (0.216); Diazinon (0.195)
				Boyreahmad	Cucumber	Endosulfan: α isomer (0.295), β isomer (0.349); Diazinon (0.669)
					Tomato	Endosulfan: α isomer (0.447), β isomer (0.435); Diazinon (0.392)
				Dena	Cucumber	Endosulfan: α isomer (0.207), β isomer (0.297); Diazinon (0.088)
					Tomato	Endosulfan: α isomer (0.131), β isomer (0.102); Diazinon (0.534)
				Total (kohkiloyeh & boyerahmad province)	Cucumber	Endosulfan: α isomer (0.255), β isomer (0.341); Diazinon (0.355)
						Tomato
Market	4 samples from each city			Yasouj	Cucumber	Endosulfan: α isomer (Undetectable), β isomer (Undetectable); Diazinon (0.121)
					Tomato	Endosulfan: α isomer (Undetectable), β isomer (0.006); Diazinon (0.030)
				Dehdasht	Cucumber	Endosulfan: α isomer (0.008), β isomer (0.016); Diazinon (0.201)
					Tomato	Endosulfan: α isomer (Undetectable), β isomer (0.011); Diazinon (0.211)
				Ghachsaran	Cucumber	Endosulfan: α isomer (0.021), β isomer (0.031); Diazinon (0.092)
					Tomato	Endosulfan: α isomer (Undetectable), β isomer (Undetectable); Diazinon (Undetectable)
(Farshad, 2001)	Wholesale market	378	GC ECD_TSD	Tehran	Cucumber	B-hch (0.2226±0.042), Linden (0.005±0.0013), Heptachlor (0.003±0.0008), Heptachlorepoxyd (0.019±0.0122), Dielrin(0.0028±0.0002), PP-DDE(0.004±0.0005), β endosulfan(0.0007±0.0004), OP-DDT(0.02±0.0007), PP-DDT(0.01±0.0215), Parathion(0.0364±0.109), Phamthion (0), Diazinon (0), Malathion (0.136±0.1129), Chloroprimumphos (0), Phirimiphos (0.085±0.0525), Phenirtathion (0.059±0.127)

Discussion

In this systematic review, 24 studies were included reporting pesticide residues in Iranian fruits and vegetables. The percentage of Iranian fruits and vegetables contaminated with pesticides exceeding the MRL was less than 10% in most studies. Samples of contaminated goods were collected mainly from cultivated areas such as fields, orchards, or greenhouses. The amount of pesticides decreased significantly after a few days; therefore, the residual toxins were less before consumption by people.

Pesticide residues in fruits and vegetables have been reported higher in Ghana (Donkor *et al.*, 2016) compared to Iran. However, the pesticides reported in this study were somewhat different from Iranian studies and the sampling sites were not mentioned. The percentage of foods contaminated with pesticides above the MRL was close to the present study although the pesticides examined were slightly different (Donkor *et al.*, 2016). Overall, pesticide residue control appears to be a serious problem in most developing countries although the extent of the problem can vary slightly. There is now enough evidence to claim misuse or even overuse of pesticides in most developing countries, mostly due to a lack of education (Ecobichon, 2001).

It is generally believed that the use of pesticides is inevitable (Carvalho, 2006) to ensure food security worldwide. On the other hand, the WHO has reported that in developing countries, 37,000 cases of cancer are linked to pesticide use each year (Tudi *et al.*, 2021). In addition, the FAO reported that three million people worldwide are poisoned by pesticides every year, and 200,000 people die each year. The worst part is that the majority of them are from developing countries (Watts, 2010). Another important problem is the use of banned pesticides in developing countries (Tariq *et al.*, 2007). Therefore, it is clear that pesticide residues must be monitored to control side effects.

In fact, pesticide residues in food have not been systematically reported in Iran. Public institutions do not systematically report residues, and only a

few articles were published by academic societies. Considering the current scenario, there is an urgent need to facilitate reliable and continuous measurements of toxic residues in Iranian food. What is more important is to create systematic training of farmers to use appropriate pesticides through effective methods such as social marketing. The authors strongly recommend that public awareness of the safe use of pesticides should be raised through mass media and social media. The control of pesticide residues based on the Codex Alimentarius is inevitable for Iran, not only because of the health of citizens but also because of the possibility of exporting food.

Conclusion

This systematic review on pesticide residues in Iran found limited studies, indicating a paucity of data. In most studies, less than 10% of Iranian fruits and vegetables were contaminated with toxins above the acceptable limits. However, at this time, we cannot make an accurate claim about pesticide residues in Iranian food, and further studies are required.

Acknowledgment

There is no acknowledgment.

Authors' contributions

Toorang F designed the study and did the primary search. Toorang F and Sasanfar B performed study selection and data extraction. Eskandari S and Pouraram H consulted the whole study. All authors read and approved the paper.

Conflict of interest

None of the authors declared any conflict of interest.

Funding

None

References

- Akhlaghi H, Motavalizadehkakhky A & Emamiyan R 2013. Determination of diazinon in fruits from northeast of Iran using the QuEChERS sample preparation method and GC/MS. *Asian journal of chemistry*. **25** (3): 1727.

- Alavanja MC, Hoppin JA & Kamel F** 2004. Health effects of chronic pesticide exposure: cancer and neurotoxicity. *Annual review of public health*. **25**: 155.
- Ardakani AS, morovati m & entesari m** 2012. Residue of endosulfan and diazinon pesticides in tomato and green cucumber fields in Kohgilouye and Boyer Ahmad provinces. *Journal of genetic engineering and biosafety*. **1 (2)**.
- Asghar U, Malik M & Javed A** 2016. Pesticide exposure and human health: a review. *Journal of ecosystem & ecography open access*. **5**: 2.
- askari m, morowati m & eimani s** 2014. Determination of Diazinon residue levels on Cherry, *Cerasus avium* supplied to Tehran central fruit and vegetable market. *Genetic engineering and biosafety journal*. **2 (2)**: 119-126.
- Bayat M, Hassanzadeh-Khayyat M & Mohajeri SA** 2015. Determination of diazinon pesticide residue in tomato fruit and tomato paste by molecularly imprinted solid-phase extraction coupled with liquid chromatography analysis. *Food analytical methods*. **8 (4)**: 1034-1041.
- Behbahaninia** 2007. Chemical contamination of cucumber product with the chemical poisons pmetrozine, deltamethrin, and tetradifon in Damavand region. *Journal of plant and ecosystem*. **2 (8)**: 118-128.
- Carvalho FP** 2006. Agriculture, pesticides, food security and food safety. *Environmental science & policy*. **9 (7-8)**: 685-692.
- Donkor A, et al.** 2016. Pesticide residues in fruits and vegetables in Ghana: a review. *Environmental science and pollution research*. **23 (19)**: 18966-18987.
- Ecobichon DJ** 2001. Pesticide use in developing countries. *Toxicology*. **160 (1-3)**: 27-33.
- Farshad A** 2001. Evaluation of Chlorine and Phosphorus Pesticide Residual in Cucumber in Tehran Fruit and Vegetable Center Market and Its Health Hazards Effects. *Internal medicine today*. **6 (14)**: 50-59.
- Ganjeizadeh Rohani F, Mahdavi V & Aminaei MM** 2014. Investigation on diazinon and oxydemeton-methyl residues in cucumbers grown in Kerman greenhouses. *Environmental monitoring and assessment*. **186 (7)**: 3995-3999.
- Golepoor M, et al.** 2014. Assessment of Organophosphorus Residues together in Strawberry Produced in Mazandaran, Iran. *Journal of Mazandaran University of Medical Sciences*. **23 (109)**: 93-102.
- Hadian Z & Azizi M** 2006. Pesticide Residues In Vegetables Marketed In The Main Wholesale Fruit And Vegetable Market In Tehran As Determined By Gas Chromatography/Mass Spectrometry, 2005. *Journal of nutrition sciences & food technology*. **1 (2)**: 13-20.
- Hadian Z & Azizi mh** 2008. Evaluation of the residual amount of pesticides by gas chromatography-mass spectrometry method in some vegetables offered in the main fruit and vegetable square of Tehran 2014. *Iranian journal of nutrition sciences & food technology*. **1 (2)**: 13-20.
- Hadian Z, Azzi MH & Ferdousi R** 2006. Determination of chlorinated pesticide residues in vegetables by gas chromatography/mass spectrometry. *Journal of food science and technology*. **3 (8)**: 67-74.
- Hagian Shahri M, Sonei A, Zohour E, Khoshbazzm R & Tagbakhsh MR** 2014. Investigation on residue of pesticides in some horticultural crops with gas chromatography method (GC/MS) in Khorassan Razavi Province. *Journal of applied research in plant protection*. **3 (2)**: 93-106.
- Herrman JI** 1993. The role of the World Health Organization in the evaluation of pesticides. *Regulatory toxicology and pharmacology*. **17 (3)**: 282-286.
- Jafari A, et al.** 2012. Monitoring dithiocarbamate fungicide residues in greenhouse and non-greenhouse tomatoes in Iran by HPLC-UV. *Food additives & contaminants: Part B*. **5 (2)**: 87-92.
- Jahanmard E, Ansari F & Feizi M** 2016. Evaluation of Quechers sample preparation and GC mass spectrometry method for the determination of 15 pesticide residues in tomatoes used in salad production plants. *Iranian journal of public health*. **45 (2)**: 230.

- Khak MT, et al.** 2016. Determining the residual cypermethrin, permethrin, indoxacarb and mancozeb in tomato produced in bushehr province farms. *International journal of medical research & health sciences*. **5 (5)**: 210-217.
- Khaniki J, Fadaei A, Sadeghi M & Mardani G** 2011. Study of Oxydimeton methyl residues in cucumber & tomato grown in some of greenhouses of Chaharmahal va Bachtari province by HPLC method. *Journal of Shahrekord University of Medical Sciences*. **13 (4)**: 9-17.
- Kim K-H, Kabir E & Jahan SA** 2017. Exposure to pesticides and the associated human health effects. *Science of the total environment*. **575**: 525-535.
- Leili M, et al.** 2016. Determination of pesticides residues in cucumbers grown in greenhouse and the effect of some procedures on their residues. *Iranian journal of public health*. **45 (11)**: 1481.
- Mayne ST, Playdon MC & Rock CL** 2016. Diet, nutrition, and cancer: past, present and future. *Nature reviews clinical oncology*. **13 (8)**: 504-515.
- Mina M & Maryam F** 2012. Determining the residual amount of diazinon and chlorpyrifos in Golden and Red apple varieties of the region. *Ecology* **38 (62)**: 111-116.
- Mohammadi S & Imani S** 2012. Deltamethrin and chlorpyrifos residue determination on greenhouse tomato in Karaj by Solid Phase Extraction. *Plant protection journal*. **4 (1)**.
- Pirsaheb M, Fattahi N, Rahimi R, Sharafi K & Ghaffari HR** 2017. Evaluation of abamectin, diazinon and chlorpyrifos pesticide residues in apple product of Mahabad region gardens: Iran in 2014. *Food chemistry*. **231**: 148-155.
- Popp J, Pető K & Nagy J** 2013. Pesticide productivity and food security. A review. *Agronomy for sustainable development*. **33 (1)**: 243-255.
- Shokerzadeh M, Vahedi H & Shabankhani B** 2006. Investigation and Measurement of Pesticidal Residues Benomil and Mancozeb in Cucumber Produced in Mazandaran. *Journal of Shahid Sadoughi University of Medical Sciences*. **13 (5)**: 59-64.
- Shokrzadeh M, Karami M, Jafari Valoujaei M & Zamani Renani A** 2013. Measuring Diazinon residue in Thompson orange. *Journal of Mazandaran University of Medical Sciences*. **23 (105)**: 91-99.
- Sobhanardakani S, Sadri S & Jameh Bozorgi S** 2014. Evaluation of organophosphorus pesticide diazinon residue in greenhouse crops using spectrophotometry (case study: mushroom). *Food hygiene*. **3 (4 (12))**: 73-80.
- Sobhanardakani S, Younesian M & Jameh Bozorgi S** 2016. Evaluation of organophosphorus pesticide diazinon residues in greenhouse crops (Case study: Zucchini). *Journal of environmental science and technology*. **18 (3)**: 141-148.
- Tariq MI, Afzal S, Hussain I & Sultana N** 2007. Pesticides exposure in Pakistan: a review. *Environment international*. **33 (8)**: 1107-1122.
- Tudi M, et al.** 2021. Agriculture development, pesticide application and its impact on the environment. *International journal of environmental research and public health*. **18 (3)**: 1112.
- Vaidyanathan G** 2021. What humanity should eat to stay healthy and save the planet. *Nature*. **600 (7887)**: 22-25.
- Watts M** 2010. Pesticides: Sowing poison, growing hunger, reaping sorrow. Pesticide Action Network Asia and the Pacific.
- World Health Organization** 2019. Healthy diet. World Health Organization. Regional Office for the Eastern Mediterranean.